UBLISHED BY THE
NTOMOLOGICAL
OCIETY OF AMERICA

JS ISSN 0022 0493
EENAI 96
2003)

EDITOR-IN-CHIEF
John T. Trumble

SUBJECT EDITORS
Arthur G. Appel
Peter W. Atkinson
Barbara J. Bentz
Michael Brewer
Jacques Brodeur
G. David Buntin
Ring T. Cardé
Robert R. Granados
William D. Hutchison
Michael E. Irwin
Murray B. Isman
Gail E. Kampmeier
Marc Klowden
Timothy J. Lysyk
Lisa G. Neven
Thomas W. Phillips
Stuart R. Reitz
Stanley S. Schneider
David J. Schuster
Anthony M. Shelton
Michael J. Stout
Julie L. Todd
Nikolai A. M. Van Beek
Robert C. Venette
James F. Walgenbach
Jian-Zhou Zhao
Kun Yan Zhu

Journal of Economic Entomology

Index
Volume 96
2003

Journal of Economic Entomology

Volume 96

EDITOR-IN-CHIEF

JOHN T. TRUMBLE
Riverside, CA

SUBJECT EDITORS

ARTHUR G. APPEL
Auburn, AL

PETER W. ATKINSON
Riverside, CA

BARBARA J. BENTZ
Logan, UT

MICHAEL BREWER
East Lansing, MI

JACQUES BRODEUR
Ste Foy, PQ, Canada

G. DAVID BUNTIN
Griffin, GA

RING T. CARDÉ
Riverside, CA

ROBERT R. GRANADOS
Ithaca, NY

WILLIAM D. HUTCHISON
Saint Paul, MN

MICHAEL E. IRWIN
Urbana, IL

MURRAY B. ISMAN
Vancouver, BC, Canada

GAIL E. KAMPMEIER
Urbana, IL

MARC KLOWDEN
Moscow, ID

TIMOTHY J. LYSYK
Lethbridge, AB, Canada

LISA G. NEVEN
Wapato, WA

THOMAS W. PHILLIPS
Stillwater, OK

STUART R. REITZ
Tallahassee, FL

STANLEY S. SCHNEIDER
Charlotte, NC

DAVID J. SCHUSTER
Bradenton, FL

ANTHONY M. SHELTON
Geneva, NY

MICHAEL J. STOUT
Baton Rouge, LA

JULIE L. TODD
Ames, IA

NIKOLAI A. M. VAN BEEK
College Park, MD

ROBERT C. VENETTE
St. Paul, MN

JAMES F. WALGENBACH
Fletcher, NC

JIAN-ZHOU ZHAO
Geneva, NY

KUN YAN ZHU
Manhattan, KS

ASSOCIATE EDITORS

PETER FOLLETT
Hilo, HI

GUY HALLMAN
Weslaco, TX

WILLIAM J. MOAR
Auburn, AL

JAQUELINE L. ROBERTSON
Petaluma, CA

BOOK REVIEWS EDITOR

ABELARDO C. MONCAYO
Ada, OH

ESA STAFF

ALAN KAHAN
Director of Communications

LATHIFAH M.-IBRAHIM
Issue Manager (Cadmus)

EDITORIAL BOARD

WILLIAM D. HUTCHISON, *Chair*
St. Paul, MN
Section F, 2003

RICHARD W. MANKIN
Gainesville, FL
Section B, 2005

TIMOTHY LYSYK
Lethbridge, AB, Canada
Section D, 2006

ALLEN E. KNUTSON
Dallas, TX
Section E, 2004

GOVERNING BOARD

Z B MAYO, Jr.
Lincoln, NE
President

KEVIN L. STEFFEY
Urbana, IL
Vice President

MICHAEL A. IVIE
Bozeman, MT
Vice President-Elect

J. E. MCPHERSON
Carbondale, IL
Past-President

LINDA J. MASON
West Lafayette, IN
Secretary-Treasurer

JAQUELINE MILLER
Sarasota, FL
Section A

ELIZABETH F. BECKEMEYER
Winder, GA
Section B

LAWRENCE A. LACEY
Wapato, WA
Section C

EDWARD T. SCHMIDTMANN
Laramie, WY
Section D

MARLIN E. RICE
Ames, IA
Section E

SHRIPAT T. KAMBLE
Lincoln, NE
Section F

WAYNE A. GARDNER
Griffin, GA
Southeastern Branch

BRIAN A. NAULT
Geneva, NY
Eastern Branch

SCOTT H. HUTCHINS
Indianapolis, IN
North Central Branch

FRANK E. GILSTRAP
College Station, TX
Southwestern Branch

JOHN D. STARK
Puyallup, WA
Pacific Branch

PAULA G. LETTICE
Lanham, MD
Executive Director

© 2003, Entomological Society of America

The *Journal of Economic Entomology* (ISSN 0022 0493) is published bimonthly (February, April, June, August, October, and December) by the Entomological Society of America, 9301 Annapolis Road, Lanham, MD 20706–3115; (301) 731–4535; FAX (301) 731–4538. The journal is printed by Cadmus Journal Services, 2905 Byrdhill Road, Richmond, VA 23228. Periodicals postage is paid at Lanham, MD, and at additional mailing offices.

Postmaster: Send change of address forms to *Journal of Economic Entomology*, ESA, 9301 Annapolis Road, Lanham, MD 20706–3115.

Inquiries regarding content, subscriptions, advertising, availability of back issues, change of address, general business matters, and manuscripts should be sent to the Publications Office, ESA, 9301 Annapolis Road, Lanham, MD 20706–3115. For information about the preparation of manuscripts, authors are urged to consult "Information for Contributors," which is printed on the inside back cover of most issues, or "Publish with ESA" located on the ESA website at http://www.entsoc.org.

Articles report the results of research only. Mention of a pesticide or a commercial or proprietary product does not constitute an endorsement or recommendation for its use. Registered names and trademarks, etc., used in this publication, even without specific indication thereof, are not to be considered unprotected by law. All articles and items published in the journal are signed and, unless otherwise stated, reflect the individual views of the authors and not necessarily the official poiats of view held by ESA. ESA assumes no responsibility for the completeness of the articles or other items or the accuracy of the conclusions reached therein.

Photocopies. Permission to photocopy journal articles for internal or personal use is granted by the Entomological Society of America provided that a fee of $4.00 per copy is paid directly to the Copyright Clearance Center, 222 Rosewood Drive, Danvers, MA 01923. Individuals may make a single copy for personal use without charge. This consent does not extend to other kinds of copying, such as that for general distribution, for advertising or promotional purposes, for creating new collective works, or for resale.

Subscriptions. 2003 prices: members, free online, $15 print; nonmember individuals, $114 print or online; institutions, $243 print or online. Subscribe to both print and online editions for 25% off the combined rate. To the above add $23 for shipping outside the continental United States ($42 for Airmail). Send payment to Entomological Society of America, c/o First Union Bank, Lockbox #4690, P.O. Box 85080, Richmond, VA 23285–4690. Inquiries should be directed to ESA, 9301 Annapolis Road, Lanham, MD 20706–3115; Phone: 301–731–4535; Fax: 301–731–4538; or pubs@entsoc.org.

Back Issues. For prices and availability contact the ESA Office at the address above. Back volumes are available in microfilm from University Microfilms International, 300 North Zeeb Road, Ann Arbor, MI 48106.

CUMULATIVE AUTHOR INDEX
Journal of Economic Entomology
Volume 96, 2003

A

Ahmad, M., 875
Ahn, Y.-J., 1479
Akhurst, R. J., 1290
Alauzet, C., 1473
Albajes, R., 1805
Alexander, P. J., 543
Ali, M. A., 259
All, J. N., 748
Allee, L. L., 1420
Allen, R. F., 996
Alm, S. R., 453
Almeida, R. P. P., 264
Aluja, M., 334, 1108, 1693
Andersen, S. B., 202, 1922
Anderson, P. L., 361
Antony, B., 322
Appel, A. G., 863
Appleby, J. H., 489
Arif, M. I., 875
Armenta, R., 649
Armstrong, J. S., 352, 1426
Arthur, F. H., 503, 510
Ashley, R., 420
Auger, M., 768
Austin, J. W., 1514
Averill, A. L., 407

B

Ban, P., 1526
Barbara, K. A., 584
Barchia, I. M., 983
Barnard, D. R., 1559
Barrett, B. A., 623
Barrigossi, J. A. F., 1160
Barrington, A., 636
Barry, J. D., 615, 1405, 1485
Basso, A., 1237
Bastos, C. S., 188
Bauce, E., 280
Baxendale, F. P., 475, 1942
Beard, C., 1290
Begna, S. H., 1193
Behle, R. W., 1066
Bellemare, J., 1221
Bellotti, A. C., 1905, 1915
Benson, E. P., 94
Berg, R. K., 1792
Bernal, C. C., 1935
Bernal, J. S., 570, 950
Berni, J., 662
Bikoba, V., 246
Bird, L. J., 1290
Blackburn, M., 441
Bläske, V.-U., 1267
Bloem, S., 1724
Blua, M. J., 1369
Boetel, M. A., 714, 1426
Bonierbale, M., 1905
Borgemeister, C., 817
Borys, R. R., 31
Bostanian, N. J., 1221
Boucher, T. J., 420
Bradley, J. R., 137
Brandhorst-Hubbard, J. L., 1704, 1770
Bridges, W. C., Jr., 755
Brooks, S. E., 1259
Brown, G. C., 21
Brown, M. D., 1786
Brown-Guedira, G. L., 1329
Bueno, J. M., 1577
Bughrara, S. S., 1898
Burd, A. D., 137
Burkness, E. C., 1230
Bush, P., 748
Butin, E., 763
Butrón, A., 1563
Byrne, F. J., 1849

C

Caballero, P., 649, 1668
Cadogan, B. L., 388
Calderwood, W., 420
Calvin, D. D., 413
Campbell, J. B., 1016
Campbell, J. F., 1617, 1958
Capinera, J. L., 584
Cappaert, D., 1898
Cardona, C., 1577
Carlson, R. B., 1426
Carpenter, A., 1119
Carpenter, J. E., 1724
Carrière, Y., 1031, 1626
Cartea, M. E., 143
Carvalho, L. A., 1719
Castaño, O., 1915
Castillejos, V., 649
Catangui, M. A., 957
Cave, R. D., 649
Chander, G. J., 246
Chandler, B. W., 826
Chandler, L. D., 131, 1855
Chapin, J. W., 1142
Chapman, J. W., 649
Charlet, L. D., 706
Chen, W., 1230
Chen, Z., 377
Chilcutt, C. F., 1174
Chiroleu, F., 231
Chitkowski, R. L., 755
Choi, B.-R., 1479
Choi, W.-I., 1479
Chu, C.-C., 543
Cilas, C., 1473
Cisneros, J., 649
Clancy, K. M., 377, 783
Clarke-Harris, D., 798
Cohen, M. B., 1935
Conant, M. M., 287
Cone, W. W., 287
Cook, S. P., 396, 1662
Credland, P. F., 489
Cristofaro, M., 1719
Crocker, R. L., 1704, 1770
Croft, B. J., 148
Crowder, D. W., 1872
Crummey, H., 35
Cruz-López, L., 1126
Cuperus, G. W., 1585

D

Daglish, G. J., 525
Dahlsten, D. L., 1455
Davey, R. B., 1608
Davis, F. M., 12
Dawson, C. G., 453
Dean, B. B., 1510
Deaton, R., 1322
Decazy, B., 1473
Dedej, S., 1215
Dejean, A., 251
Delaplane, K. S., 1215, 1647
Delettre, Y. R., 126
Denholm, I., 875
Dennehy, T. J., 1031, 1307
De Oliveira, D., 401
Derouen, S. M., 1612
Dhileepan, K., 148
Dively, G. P., 914, 925, 931
Djiéto-Lordon, C., 251
Dong, B., 1798
Dorn, S., 1577
Dregseth, R. J., 1426
Dufour, B., 1473
Duncan, L. W., 835
Dunkel, F. V., 220, 1045
Dupont, A., 768
Duran, D. P., 599
Durgy, R., 420
Durier, V., 1254, 1832
Duthie, J., 64
Duyck, P. F., 231
Dwivedi, S. L., 1886
Dynok, S. A., 826

E
Edelson, J. V., 64
Elkinton, J., 763
Ellersieck, M. R., 599
Elliott, N. C., 975, 1585
Ellis, J. D., Jr., 1647
Ellis, M. D., 259
Ellsbury, M. M., 599, 1153
Elzen, P. J., 1647
Ems-Wilson, J., 1275
English-Loeb, G., 433
Equihua, M., 334
Evenson, P. D., 714

F
Fabre, A., 251
Fabre, F., 231
Farone, W., 636
Fei, H., 461
Fielding, D. J., 1193
Fields, P. G., 1005
Fisher, J. R., 969
Fitzgerald, C. J., 1828
Flanders, K. L., 1704, 1770
Fleischer, S. J., 798
Flinn, P. W., 996
Foil, L. D., 1612
Fondren, K. M., 1760
Forschler, B. T., 1267
Fournier, D., 468
Franke, D. E., 1612
Frei, A., 1577
Fries, I., 1641
Fuller, B. W., 714
Funderburk, J. E., 1201
Fuzy, E. M., 1076

G
Gable, R. H., 699
Gaff, N., 730
Gao, X., 1838
García, L. F., 649
Gates, D., 615
Geden, C. J., 1602
George, J. E., 1608
Giles, K. L., 975, 1184, 1585
Gill, B. S., 1329
Gilmour, A. R., 983
Gmitter, F. G., Jr., 835
Gobbi, A., 1054
Gonsiska, P., 81
González-Hernández, H., 1126
Gore, J., 699
Gore, J. C., 1362
Gosselin, D. C., 1750
Gould, F., 137, 156, 173
Goulet, H., 1221
Goulson, D., 649, 1668
Grace, J. K., 456
Grafton-Cardwell, E. E., 1388
Gray, M. E., 737, 1872
Greenplate, J. T., 1322
Groden, E., 1653
Groves, R. L., 1
Gu, H., 1577, 1798
Gu, P., 1388
Guillén, M., 577
Guo, Y., 909, 1322, 1838
Guse, C. A., 1872
Gut, L. J., 58

H
Hagood, S., 348
Hagstrum, D. W., 996
Haile, F. J., 1168
Hallman, G. J., 1399
Hammack, L., 1153
Hammon, R. W., 352
Han, O.-K., 207
Hanks, L. M., 370
Hara, A. H., 1334
Hardin, M. R., 1675
Harman, D. M., 53
Harvey, F. E., 1750
Harvey, T. L., 1329
Havill, N., 763
Hawley, C. J., 214
Hawthorne, D. J., 1021
Hayman, D. L., 1245
Hazzard, R. V., 1653
He, K., 935
He, X., 1119
Hébert, C., 768
Hein, G. L., 1160
Held, D. W., 81
Hellmich, R. L., 361
Henderson, G., 461
Heng-Moss, T. M., 475, 1942
Henneberry, T. J., 322, 543
Hepburn, R., 1647
Heraty, J. M., 577
Herbert, D. A., 914, 925
Hernandez, E., 1132
Hernández-Ortiz, V., 1693
Herron, G. A., 983
Hertel, H., 1267
Hesler, L. S., 1792
Hewa-Kapuge, S., 1083
Hibbard, B. E., 599
Higley, L. G., 188, 194, 1160, 1168
Hiller, L. K., 1510
Hinton, J. L., 1352
Hoelscher, J. A., 623
Hoffmann, A. A., 1083
Hoffmann, M. P., 361
Hofmeyr, J. H., 1724
Holbrook, G. L., 1548
Holler, T., 1719
Holloway, G., 35
Holsten, E. H., 31
Hooper-Bùi, L. M., 98
Horner, T. A., 914, 925, 931
Hou, X., 1005
Hoy, C. W., 1494
Hoy, M. A., 564, 1685
Hunt, T. E., 536, 1168, 1750
Hunter, W. B., 811
Hurlbert, S. H., 533
Hutchison, W. D., 1230, 1500, 1814

I
Ibrahim, S. A., 461
Inglis, G. D., 12, 272

J
Jackson, C. G., 543
Jacobsen, C. M., 1334
James, R. R., 25, 1091
James, W., 1290
Jang, E. B., 1719
Jaramillo, G. I., 856
Jaronski, S. T., 1045
Jespersen, J. B., 1300
Jiménez-Pérez, A., 592
Jobin, L., 768
Johnson, D. W., 21
Johnson, J. A., 519, 1952
Johnson, W. C., 690
Jones, D. B., 975, 1585
Jones, S. C., 1520
Jones, W. A., 1091
Joyce, A. L., 370

K
Kaga, A., 207
Kanga, L. H. B., 1091
Karut, K., 543
Kashiwaba, K., 207
Kay, B. H., 1786
Keena, M. A., 43
Kemp, K., 1958
Kenne, M., 251
Kennedy, G. G., 1, 156, 173
Kerns, D. L., 805
Ketunuti, U., 292
Khanuja, S. P. S., 990
Kindler, S. D., 975, 1585
Knodel, J. J., 706
Koch, G. W., 783
Koehler, P. G., 1259
Kolb, T. E., 377
Komuta, K., 555
Koppenhöfer, A. M., 1076
Kravchenko, A. N., 1898
Kristensen, M., 1300
Kroening, M. K., 352
Kumar, K. A., 1039
Kumar, S., 990, 1594

L
Lage, J., 202, 1922
Lapointe, S. L., 811
Lasa, R., 1668
Lascano, R. J., 1174
Lasnier, J., 1221

Lawrence, A. M., 12, 272
Lawson, A. B., 1455
Lax, A. R., 117
Lee, E.-H., 1479
Lee, K., 1942
Legaspi, J. C., 950
Legendre, B. L., 1929
Lei, T. T., 730
Leng, X., 1283
Leonard, B. R., 699
Leonard, D. S., 1448
Lesage, L., 1221
Levine, E., 1872
Leyva, J., 1126
Li, A. Y., 1307
Li, S. Y., 35
Li, W., 1283
Liang, G., 909
Liburd, O. E., 1821
Liedo, P., 1132
Lisha, V. S., 322
Liu, T.-X., 1230
Liu, Y. B., 805, 1100
Llanderal-Cazares, C., 1126
Lohmeyer, K. H., 748
Long, B. B., 407
López, C., 1805
Losey, J. E., 1420
Lowery, D. T., 879, 1864
Luck, R. F., 577, 1375
Lux, S., 1719
Lv, N., 1322

M

Mabry, T. R., 1738
Macedo, P. A., 259
Macedo, T. B., 188, 194, 475
Macías-Ordóñez, R., 1693
MacKay, A. J., 1612
MacKenzie, K. E., 1245
Madhavan, S., 188
Magallanes, R., 649
Mahroof, R., 1345
Malik, R., 1329
Malvar, R. A., 1563
Mangan, R. L., 341
Mankin, R. W., 1704, 1770
Manso, F., 1237
Marçon, P. C. R. G., 1016
Margolies, D. C., 482
Markwell, J. P., 475
Martin, P. A. W., 441
Martin, T., 468
Martínez, A. M., 649
Matsuda, Y., 555
Mauffette, Y., 1221
Mayama, S., 555
Mayo, J. H., 1448
McCoy, C. W., 835
McCullough, D. G., 1760
McDougall, S., 1083
McGovern, W. L., 311
McGuire, M. R., 1066
McInnis, D., 1132
McInnis, D. O., 615
McKenzie, C. L., 811
McNew, R. W., 892
McPheron, B. A., 1693, 1711
McPherson, R. M., 645, 690
Meikle, W. G., 533
Meinke, L. J., 131, 1750, 1855
Meissner, H. E., 850
Meixner, M. D., 1711
Mello, K., 1849
Meng, F., 1838
Menon, A., 1345
Millar, J. G., 370
Miller, G. T., 1340
Miller, H. R., 673
Miller, J. A., 1608
Miller, N. W., 1405, 1485, 1780
Mills, W. A., III, 690
Mirkov, T. E., 950
Mitcham, E. J., 246
Mitchell, P. D., 1872
Moar, W. J., 137
Momol, M. T., 1201
Monetti, L., 143
Montgomery, M., 763
Mony, R., 251
Moon, R. D., 1352
Moore, C. B., 1548
Morales-Ramos, J. A., 106
Moreau, G., 280
Morgan, D. J. W., 1369
Mori, H., 555
Morin, S., 1031
Morse, J. G., 615, 629, 1375, 1405
Moser, B., 98
Moses, A. L., 1719
Moyer, J. W., 1
Mullen, M. A., 1617
Mullinix, B. G., Jr., 690
Mulrooney, J. E., 311
Muñoz, D., 1668
Murillo, R., 1668
Musser, F. R., 71

N

Nansen, C., 1617
Nault, B. A., 1843
Nava, D. E., 609
Nayak, M. K., 525
Nechols, J. R., 482
Nemeth, M. A., 413
Ngollo, E. D., 1653
Ni, X., 194, 475
Nichols, R. L., 1307
Nigg, H. N., 835
Niimi, B., 1750
Nogawa, S., 555
Nonomura, T., 555
Nowatzki, T. M., 1750

O

O'Brien, G. M., 1905
Obrycki, J. J., 1184
Ochou, O. G., 468
Oehler, D. D., 1608
Oi, F. M., 1259
Olson, S. M., 1201
Onstad, D. W., 1872
Opit, G. P., 482
Ordás, A., 143
Oride, L., 1485
Orivel, J., 251
Osbrink, W. L. A., 117
Ostlie, K. R., 188
Otsu, Y., 555
Ovruski, S., 1108
Owens, C. B., 1514

P

Paine, T. D., 370
Palaniswami, M. S., 322
Palermo, B. L., 783
Palmer, T., 636
Pampapathy, G., 1886
Parajulee, M. N., 239
Parimi, S., 131
Park, H.-M., 1479
Peairs, F. B., 214, 352, 673
Peck, S. L., 156, 173
Peebles, F. S., 690
Penagos, D. I., 649
Peng, Y., 1838
Perera, O. P., 885
Perez, J., 287
Pérez-Staples, D., 1693
Persad, A. B., 564
Peters, B. C., 1828
Peterson, C. J., 1275
Peterson, R. K. D., 536
Phillips, J. M., 892
Phillips, T. W., 1617
Picoaga, A., 143
Pikul, J. L., Jr., 1153
Pilcher, C. D., 941
Piñero, J., 334, 1693
Piñero, J. C., 826, 1485
Plate, J., 1843
Podgwaite, J. D., 1662
Poehling, H.-M., 817
Pons, X., 1805
Poopathi, S., 1039
Postali Parra, J. R., 609
Potter, D. A., 81, 1412
Pound, J. M., 1608
Powell, M. R., 300
Prajapati, V., 990, 1594
Pranschke, A. M., 98
Prokopy, R. J., 826, 1485
Puche, H., 88
Purcell, A. H., 264
Purdie, D. M., 1786
Puterka, G. J., 636

Putnam, S., 1750

Q

Qiu, H.-M., 1798
Qiu, X., 1283
Quesada-Allué, L. A., 662, 1237
Quick, J. S., 352
Quilici, S., 231
Quisenberry, S. S., 194, 475

R

Rabossi, A., 662
Raffa, K. F., 777
Raina, S., 1641
Rancourt, B., 401
Randolph, T. L., 214, 352, 673
Raw, A. S., 1719
Reagan, T. E., 1929
Reardon, B. J., 328
Reardon, R. C., 1662
Reay-Jones, F. P. F., 1929
Rebek, E. J., 446
Rector, B. G., 909
Reddy, L. J., 1886
Reed, C. R., 996
Reekie, E. G., 1245
Reissig, W. H., 1463
Reitz, S. R., 1201
Rendon, P., 1132
Rhainds, M., 433
Rice, M. E., 941
Rideout, T., 35
Rieske, L. K., 777
Riis, L., 1905, 1915
Riley, D. G., 1315
Riley, M. B., 94
Riordan, T. P., 1942
Rivault, C., 1254, 1832
Roberts, P. M., 748
Roberts, W., 64
Roebuck, J., 1548
Roehrdanz, R. L., 669, 1153
Roesli, R., 1958
Rogers, M. E., 1412
Rojas, J. C., 1126
Rojas, M. G., 106
Rondon, S. I., 737
Rorie, R. W., 892
Rosenkrans, C. F., Jr., 892
Rothwell, N. L., 792
Roulston, T. H., 871
Roush, R. T., 1031
Rowan, J. M., 1967
Royer, T. A., 975, 1585
Ruiz-Montiel, C., 1126
Russell, T. L., 1786
Rust, M. K., 844
Ryan, P. A., 1786
Ryckewaert, P., 231

S

Sabara, H. A., 547
Sadof, C. S., 446
Sakuratani, Y., 555
Salgado, S., 1132
Salin, C., 126
Sanson, D. W., 1612
Santiago, R., 1563
Sarzynski, E. M., 1821
Schaefer, C. W., 1367
Schal, C., 1362, 1548
Scharbach, R. D., 388
Scharf, M. E., 131, 1855
Scheffrahn, R. H., 1526
Schliserman, P., 1108
Schneider, J. C., 1433
Schneider, M. I., 1054
Schroeder, A. J., 1426
Schultz, B. B., 1653
Schweizer, H., 1375
Sciabarrasi, M., 420
Scott, W. P., 902
Seagraves, M. P., 645
Seidlecki, E., 1653
Setamou, M., 817, 950, 1929
Sharkey, M. J., 21
Sharma, H. C., 1886
Shea, P. J., 31
Shelly, T. E., 1132
Shelton, A. M., 71, 1031, 1843
Shelton, T. G., 456
Sheppard, W. S., 1711
Shields, E. J., 1420
Shimizu, H., 555
Showler, A. T., 680
Shrewsbury, P. M., 1675
Siegfried, B. D., 131, 1016, 1750, 1855
Sikorowski, P. P., 272
Silva, E. M., 1510
Silva, J. G., 1711
Silverman, J., 850, 871
Simpson, R. B., 892
Simpson, S. E., 835
Sisterson, M. S., 805, 1031
Skinner, A. C., 35
Skoda, S. R., 1016
Skovmand, B., 202, 1922
Slosser, J. E., 239
Smagghe, G., 1054
Smirle, M. J., 879, 1864
Smith, C. M., 1329, 1571
Smitley, D. R., 792, 1898
Snodgrass, G. L., 902
Soengas, P., 143
Song, Y., 935
Sonvico, A., 1237
Sotelo, J., 1563
Souto, X. C., 1563
Spangler, S. M., 413
Spencer, J. L., 1738, 1872
Spurgeon, D. W., 328, 348
Stark, J. D., 1780
Starkey, S., 1571
Stavisky, J., 1201
Steck, G. J., 1711
Steelman, C. D., 892
Steinkraus, D. C., 1602
Stone, D. M., 35
Storer, N. P., 156, 173, 1530
Straka, T. J., 1448
Stringham, S. M., 1362
Strong, C. A., 885
Stuart, R. J., 835
Su, N.-Y., 88, 1526
Subramanyam, B., 1345, 1958, 1967
Suh, C. P.-C., 348
Sullivan, M. J., 755
Suphantharika, M., 292
Swartz, H. J., 53
Szalanski, A. L., 1514

T

Tabashnik, B. E., 805, 1031
Tamez-Guerra, P., 1066
Tan, W., 1315
Tang, J., 519
Tao, F. L., 1585
Teera-Arunsiri, A., 292
Thoeming, G., 817
Thomas, D. B., 1732
Thomas, G. D., 1016
Thomas, J. S., 1142
Throne, J. E., 510, 1345
Tian, Y., 1283
Tipping, C., 246
Toews, M. D., 1967
Tollerup, K., 629
Tolmay, V., 475
Tomooka, N., 207
Tomov, B. W., 570
Tosa, Y., 555
Toscano, N. C., 1849
Toyoda, H., 555
Tran, L. C., 1485
Tripathi, A. K., 990, 1594
Turnipseed, S. G., 755

U

Ulloa-Chacón, P., 856
Umble, J. R., 969

V

Vaissayre, M., 468
Valero, K. A., 1952
Valle, J., 649
Valles, S. M., 885
Van Duyn, J. W., 137, 156, 173
Vargas, R. I., 1405, 1485, 1780
Varón, J., 334
Vaughan, D. A., 207

Vaughn, T. T., 1738
Vázquez, A., 334
Vega, S. Y., 844
Vernon, P., 126
Villalobos, E., 1132
Villavaso, E. J., 311
Vincent, C., 401, 1221
Viñuela, E., 1054
Virmani, S. S., 1935

W
Waldron, J. K., 1420
Waldstein, D. E., 58
Waldvogel, M. G., 1362, 1548
Walgenbach, J. F., 1
Walker, A. M., 1685
Walker, C. B., 352
Wallbank, B. E., 525
Wang, Q., 592, 1119
Wang, S., 519
Wang, Z., 935
Warren, K. J., 1750
Watson, D. W., 1362
Watts, V. M., 220
Way, M. O., 1929
Webb, R. E., 1662
Wegbe, K., 1473
Weiser, L. A., 1184
Weiss, M. J., 361
Wen, L., 935
Wennemann, L., 287
Wilde, G. E., 503
Williams, T., 649, 1668
Wilson, L. T., 1174
Winston, M. L., 547
Wold, S. J., 1500, 1814
Wright, L. C., 287
Wright, M. G., 361
Wu, J., 259
Wu, J.-C., 1798
Wu, K., 1322, 1838
Wyatt, W. E., 1612

X
Xue, R.-D., 1559

Y
Yang, G.-Q., 1798
Yao, Z., 935
Yearby, E. L., 1201
Yildirim, E., 1494
Yokoyama, V. Y., 1340
Young, L., 1942
Yue, B., 503

Z
Zbarsky, V., 1420
Zhang, M., 1704, 1770
Zhao, J.-Z., 1031, 1843
Zheng, Y., 53
Zhou, D., 935
Zhou, X., 1855
Zungoli, P. A., 94
Zurek, L., 1362
Zurowski, C. L., 879

CUMULATIVE SUBJECT INDEX
Journal of Economic Entomology
Volume 96, 2003

A

abamectin,
elm leaf beetle, 1455
imidacloprid, 1455
integrated pest management, 1455
Xanthogaleruca luteola, 1455
above- below-ground growth,
grasshopper density, 1193
Melanoplus sanguinipes, 1193
oilseed crop, 1193
small-grains, 1193
acephate,
chlorpyrifos, 645
lambda-cyhalothrin, 645
methomyl, 645
Solenopsis invicta, 645
toxicology, 645
Aceria tosichella,
Aegilops tauschii, 1329
plant resistance, 1329
Triticum aestivum, 1329
acyl sugar,
biopesticide, 636
biorational insecticide, 636
polyol esters, 636
sucrose octanoate, 636
surfactant, 636
Adelges tsugae,
biological control, 763
hemlock, 763
hemlock woolly adelgid, 763
Pseudoscymnus tsugae, 763
Scymnus ningshanensis, 763
adult emergence, 714, 925
corn rootworms, 714
Helicoverpa zea, 925
resistance management, 925
root injury, 714
soil insecticide rates, 714
transgenic corn, 925
Aedes,
Bacillus thuringiensis variety *israelensis*, 1786
biocontrol, 1786
Culex, 1786
Ochlerotatus, 1786
Aedes albopictus,
Anopheles quadrimaculatus, 1559
Culex nigripalpus, 1559
toxicity, 1559
Aegilops tauschii, 1329, 1571
Aceria tosichella, 1329
antibiosis, 1571
plant resistance, 1329
Schizaphis graminum, 1571
tolerance, 1571
Triticum aestivum, 1329
wheat, 1571
aesthetic injury level,
balsam fir, 1760
Fraser fir, 1760
host plant phenology, 1760
integrated pest management, 1760
Aethina tumida,
Apis mellifera, 1647
integrated pest management, 1647
modified hive entrances, 1647
screened bottom boards, 1647
Africa,
American foulbrood, 1641
Apis mellifera subsp. *scutellata*, 1641
age,
Cnephasia jactatana, 592
mating delay, 592
reproductive potential, 592
Tortricidae, 592
aggregation pheromone,
Coleoptera, 1126
Curculionidae, 1126
Scyphophorus acupunctatus, 1126
AK (south-central),
antiaggregation, 31
Bark beetles, 31
Dendroctonus rufipennis, 31
Picea sp., 31
semiochemicals, 31
aldicarb,
Cotton, 748
Frankliniella fusca, 748
precision placement, 748
Temik, 748
tobacco thrips, 748
alfalfa,
alfalfa weevil, 536
bean leaf beetle, 536
Monte Carlo simulation, 536
soybean, 536
alfalfa weevil,
alfalfa, 536
bean leaf beetle, 536
Monte Carlo simulation, 536
soybean, 536
alginate beads,
bacteriophages, 555
chitinase *SH1*, 555
Enterobacter cloacae, 555
Epilachna vigintioctopunctata, 555
Allium cepa,
honey bee behavior, 1510
honey bees, 1510
onion, 1510
pollination, 1510
Alphitobius diaperinus, 126, 1352, 1602
Beauveria bassiana, 1602
control, 126
Dermestes maculatus, 1602
field trial, 126
house fly, 1352
insect growth regulator, 126
lesser mealworm, 1352
Musca domestica, 1352
poultry, 1602
poultry house, 126
poultry pest management, 1352
pyrethroid, 126
Amaranthus spp.,
callaloo, 798
Herpetogramma bipunctalis, 798
Spodoptera spp., 798
Spoladea recurvalis, 798
American cockroach,
dog, 1259
drywood termites, 1259
subterranean termites, 1259
termite detection, 1259
termite-damaged wood, 1259
American foulbrood,
Africa, 1641
Apis mellifera subsp. *scutellata*, 1641
Anagrapha falcifera nuclear polyhedrovirus,
lignin, 1066
microencapsulation, 1066
shelf-life, 1066
spray dry, 1066
Anasa tristis,
citrullus lanatus, 64
Cucurbitaceae, 64
Hemiptera, 64
Anastrepha,
attractants, 334
human urine, 334

Tephritidae, 334
Anastrepha fraterculus, 1108, 1693
Argentina, 1108
Ceratitis capitata, 1108
citrus, 1693
host plants, 1108
host status, 1693
oviposition behavior, 1693
pest status, 1693
Tephritidae, 1108
Anastrepha fraterculus' stocks,
isolation barriers, 1237
karyotype variability, 1237
N-banding, 1237
RAPDs/PCR, 1237
Anopheles quadrimaculatus,
Aedes albopictus, 1559
Culex nigripalpus, 1559
toxicity, 1559
ant exclusion,
granular contact insecticide, 98
perimeter treatment, 98
repellency, 98
Anthonomus grandis, 348, 680
boll weevil, 348, 680
cotton, 680
kill strips, 348
pheromone traps, 348
trapping, 680
Anthonomus grandis grandis,
boll weevil, 328
colonization, 328
cotton, 328
Anthonomus musculus,
cranberry, 407
cranberry weevil, 407
plant compensation, 407
Vaccinium, 407
yield, 407
antiaggregation,
AK (south-central), 31
Bark beetles, 31
Dendroctonus rufipennis, 31
Picea sp., 31
semiochemicals, 31
antibiosis, 811, 1571, 1577, 1922, 1935
Aegilops tauschii, 1571
antifeedant, 811
antixenosis, 1577, 1922
Citrus, 811
cover crops, 811
Diaprepes abbreviatus, 811
heterobeltiosis, 1935
heterosis, 1935
host plant resistance, 1577, 1935
hybrid rice, 1935
Phaseolus vulgaris, 1577
Schizaphis graminum, 1571, 1922
synthetic hexaploid wheat, 1922
Thrips palmi, 1577
tolerance, 1571, 1922
wheat, 1571
Anticarsia gemmatalis,
Empoasca fabae, 690
genetically modified soybeans, 690
Nezara viridula, 690
Roundup Ready soybeans, 690
Spissistilus festinus, 690
antifeedant,
antibiosis, 811
Citrus, 811
cover crops, 811
Diaprepes abbreviatus, 811
antixenosis, 1577, 1922
antibiosis, 1577, 1922
host plant resistance, 1577
Phaseolus vulgaris, 1577
Schizaphis graminum, 1922
synthetic hexaploid wheat, 1922
Thrips palmi, 1577
tolerance, 1922
ants, 251, 856
biological control, 251
hemiptera, 251
insecticides, 251, 856
Tapinoma melanocephalum, 856
territoriality, 251
toxic baits, 856
Aphis glycines,
fluorescence, 188
gas exchange, 188
insect-plant relations, 188
photosynthesis, 188
Aphis gossypii, 246, 875
Frankliniella occidentalis, 246
insecticide resistance, 875
organosilicones, 246
Pakistan, 875
Platynota stultana, 246
Pseudococcus maritimus, 246
pyrethroids, 875
Tetranychus pacificus, 246
Aphis pomi,
green apple aphid, 1864
imidacloprid, 1864
mortality, 1864
toxicity, 1864
Apis mellifera, 259, 547, 1091, 1215, 1647
Aethina tumida, 1647
Apistan, 259
application methods, 1091
bee density, 1215
biological control, 1091
CheckMite+, 259
colony management, 547
greenhouse, 547
honey bee, 259, 547
integrated pest management, 1647
Metarhizium anisopliae, 1091
modified hive entrances, 1647
pollination, 547, 1215
rabbiteye blueberry, 1215
screened bottom boards, 1647
tebufenpyrad, 259
tomatoes, 547
Vaccinium ashei, 1215
varroa, 259
Varroa destructor, 1091
Apis mellifera subsp. *scutellata*,
Africa, 1641
American foulbrood, 1641
Apistan,
Apis mellifera, 259
CheckMite+, 259
honey bee, 259
tebufenpyrad, 259
varroa, 259
apple,
Grapholita molesta, 58
mating disruption, 58
microencapsulation, 58
sprayable pheromone, 58
application methodology,
entomopathogenic nematodes, 287
Heterorhabditis bacteriophora, 287
Steinernema spp., 287
Vitis, 287
application methods,
Apis mellifera, 1091
biological control, 1091
Metarhizium anisopliae, 1091
Varroa destructor, 1091
Arachis hypogaea,
Cyrtomenus ciliatus, 1142
economic injury, 1142
Pangaeus bilineatus, 1142
Sehirus cinctus, 1142
Arachis spp.,
groundnut, 1886
Helicoverpa, 1886
resistance, 1886
Spodoptera, 1886
areawide pest management,
Eoreuma loftini, 1929
host plant resistance, 1929
sugarcane, 1929
Argentina,
Anastrepha fraterculus, 1108
Ceratitis capitata, 1108
host plants, 1108
Tephritidae, 1108
Argentine ant, 844, 871
control, 844
foraging, 871

foraging range, 844
granular bait, 871
Linepithema humile, 844, 871
resurgence, 844
aromatic cedar mulch,
cultural control, 850
field study, 850
Linepithema humile, 850
nesting sites, 850
arrhenotokous,
dosage compensation, 1626
haplodiploid resistance, 1626
resistance in natural enemies, 1626
sexual dimorphism resistance, 1626
associational resistance,
interplanting, 81
pest management, 81
Popillia japonica, 81
Rosa × *hybrida*, 81
Ataenius spretulus,
carabids, 792
golf course, 792
milky disease, 792
Paenibacillus sp., 792
staphylinids, 792
attractants, 334, 826
Anastrepha, 334
Conotrachelus nenuphar, 826
human urine, 334
monitoring, 826
odor, 826
Tephritidae, 334
trap trees, 826
augmentative biological control, 1375, 1675
Citrus sinensis, 1375
cost analysis, 1675
heteronomous hyperparasitoid, 1375
integrated pest management, 1675
natural enemies, 1675
ornamentals, 1675
Australia,
bait technology, 1828
chitin synthesis inhibitor, 1828
chlorfluazuron, 1828
subterranean termites, 1828
Azadirachta indica,
neem, 817
soil application, 817
systemic effects, 817
western flower thrips, 817
azadirachtin,
Bemisia, 25
microbial control, 25
Paecilomyces, 25

B

B. thuringiensis,
biopesticides, 292
formulation, 292
spray drying, 292
suspensibility, 292
wettability, 292
Bacillus thurengiensis var. *kurstaki*,
diflubenzuron, 396
Orgyia pseudotsugata, 396
tebufenozide, 396
Bacillus thuringiensis, 137, 156, 173, 280, 388, 879, 1031, 1290, 1322, 1653, 1838
Beauveria bassiana, 1653
bollworm, 137
Bt crops, 1031
Bt resistance, 137
Chilo suppressalis, 1838
China, 1322, 1838
Choristoneura fumiferana, 280
Choristoneura rosaceana, 879
computer simulation, 156, 173
Double application, 280
efficacy, 388
experimental design, 388
fitness, 1290
gene frequency, 137
genetically modified crops, 1031
Helicoverpa armigera, 1290, 1322
Helicoverpa zea, 1653
insecticide resistance, 879
orchard pest management, 879
Ostrinia nubilalis, 1653
Pandemis limitata, 879
refuge, 1031
replication, 388
resistance, 1031, 1290
resistance management, 156, 173
spinosad, 879
spruce budworm, 280, 388
sublethal effects, 280
susceptibility, 1838
sweet corn, 1653
transgenic cotton, 1290, 1322
transgenic crops, 156, 173
Bacillus thuringiensis corn,
European corn borer, 957
permethrin, 957
South Dakota, 957
univoltine ecotype, 957
yield, 957
Bacillus thuringiensis subsp. *israelensis*,
Bengal gram, 1039
common sugar, 1039
Luria Bertani, 1039
mosquitoes, 1039
potato, 1039
Bacillus thuringiensis variety *israelensis*,
Aedes, 1786
biocontrol, 1786
Culex, 1786
Ochlerotatus, 1786
bacteriophages,
alginate beads, 555
chitinase *SH1*, 555
Enterobacter cloacae, 555
Epilachna vigintioctopunctata, 555
Bactrocera cucurbitae, 231, 1485
bait sprays, 1485
concentration., 231
food attractants, 231
GF-120 fruit fly bait, 1485
protein hydrolysates, 231
spinosad, 1485
baculovirus,
crop phenology, 649
maize, 649
natural enemies, 649
synthetic pesticides, 649
bait, 311, 441, 1520
boll weevil, 311
corn rootworm, 441
cucurbitacin E-glycoside, 441
Diabrotica undecimpunctata howardi, 441
insect control, 441
malathion, 311
placement, 1520
Reticulitermes flavipes, 1520
Sentricon, 1520
subterranean termite control, 1520
toxicity, 311
bait contamination,
Solenopsis invicta, 94
bait sampling,
garden symphylan, 969
Scutigerella immaculata, 969
soil ecology, 969
soil sampling, 969
symphylid, 969
bait sprays,
Bactrocera cucurbitae, 1485
GF-120 fruit fly bait, 1485
spinosad, 1485
bait technology,
Australia, 1828
chitin synthesis inhibitor, 1828
chlorfluazuron, 1828
subterranean termites, 1828
balsam fir,
aesthetic injury level, 1760
Fraser fir, 1760
host plant phenology, 1760
integrated pest management, 1760

Bark beetles,
AK (south-central), 31
antiaggregation, 31
Dendroctonus rufipennis, 31
Picea sp., 31
semiochemicals, 31
bean leaf beetle,
alfalfa, 536
alfalfa weevil, 536
Monte Carlo simulation, 536
soybean, 536
Beauveria bassiana, 1602, 1653
Alphitobius diaperinus, 1602
Bacillus thuringiensis, 1653
Dermestes maculatus, 1602
Helicoverpa zea, 1653
Ostrinia nubilalis, 1653
poultry, 1602
sweet corn, 1653
bee density,
Apis mellifera, 1215
pollination, 1215
rabbiteye blueberry, 1215
Vaccinium ashei, 1215
beef cattle,
heifers, 1612
Hematobia irritans, 1612
reproduction, 1612
weight gain, 1612
beetles,
chlorpyrifos-methyl, 525
psocids, 525
sorghum, 525
synergized bifenthrin, 525
behavior,
integrated pest management, 401
Lygus lineolaris, 401
physical control method, 401
strawberry, 401
vacuuming, 401
Bemisia, 25, 1307
azadirachtin, 25
insect growth regulator (IGR), 1307
microbial control, 25
Paecilomyces, 25
pyriproxyfen, 1307
resistance, 1307
whitefly, 1307
Bemisia argentifolii,
bifenthrin resistance, 1315
host plant, 1315
inheritance, 1315
Bemisia tabaci, 322, 543
biotype, 322
Encarsia, 543
Eretmocerus, 543
esterase, 322
LED-plastic cup trap, 543
virus, 322
yellow sticky card trap, 543
beneficial,
Hymenoptera, 1083
integrated pest management, 1083
pesticide, 1083
Trichogramma, 1083
Bengal gram,
Bacillus thuringiensis subsp. *israelensis*, 1039
common sugar, 1039
Luria Bertani, 1039
mosquitoes, 1039
potato, 1039
Beta vulgaris,
cover crop, 1426
integrated control, 1426
soil insecticide, 1426
Tetanops myopaeformis, 1426
bifenthrin resistance,
Bemisia argentifolii, 1315
host plant, 1315
inheritance, 1315
bioassay, 220, 377, 983, 1016
budburst phenology, 377
Choristoneura occidentalis, 377
defoliation, 377
dry weight loss, 220
extra binomial variation, 983
generalized linear mixed model, 983
host plant resistance, 220
insecticide resistance, 1016
kernel damage, 220
methoxychlor, 1016
Musca domestica, 1016
permethrin, 1016
probit analysis, 983
Pseudotsuga menziesii, 377
resistance mechanisms, 377
Rhyzopertha dominica, 220
spray oils, 983
stirofos, 1016
stored-products, 220
bioavailability,
isoborneol, 1267
persistence, 1267
Reticulitermes spp., 1267
soil treatments, 1267
biocontrol,
Aedes, 1786
Bacillus thuringiensis variety *israelensis*, 1786
Culex, 1786
Ochlerotatus, 1786
bioinvasion,
Ceratitis capitata, 1711
microsatellites, 1711
mtDNA, 1711
polymerase chain reaction-restriction fragment length polymorphism, 1711
biological control, 21, 251, 370, 446, 570, 706, 763, 1045, 1091
Adelges tsugae, 763
ants, 251
Apis mellifera, 1091
application methods, 1091
biorational pesticide use, 446
cultural control, 706
Eoreuma loftini, 570
fitness, 570
Galanthus nivalis agglutinin, 570
hemiptera, 251
hemlock, 763
hemlock woolly adelgid, 763
insecta, 370
insecticides, 251
integrated pest management (IPM), 21
life table, 21
Melanoplus sanguinipes, 1045
Metarhizium anisopliae, 1091
mycoinsecticide, 1045
Myiopharus macellus, 706
nontarget effects, 570
nontarget organism, 1045
nutrition, 370
parasitoids, 446
Phoracantha recurva, 370
Phoracantha semipunctata, 370
pollen, 370
predators, 1045
Pseudoscymnus tsugae, 763
scale insects, 446
Scymnus ningshanensis, 763
stored product insects, 1045
sunflower beetle, 706
systemic insecticides, 446
territoriality, 251
Varroa destructor, 1091
Zygogramma exclamationis, 706
biopesticide,
acyl sugar, 636
biorational insecticide, 636
polyol esters, 636
sucrose octanoate, 636
surfactant, 636
biopesticides,
B. thuringiensis, 292
formulation, 292
spray drying, 292
suspensibility, 292
wettability, 292
biorational insecticide,
acyl sugar, 636
biopesticide, 636
polyol esters, 636
sucrose octanoate, 636
surfactant, 636

biorational pesticide use,
biological control, 446
parasitoids, 446
scale insects, 446
systemic insecticides, 446
biotype,
Bemisia tabaci, 322
esterase, 322
virus, 322
Blattella germanica, 863, 1254, 1548, 1832
cross-resistance, 1548
cyclodienes, 1548
Ebeling choice box, 863
edge effect, 1832
efficiency, 1254
exploratory behavior, 1832
field performance, 863
Fipronil, 1548
fipronil gel bait, 1254
fragmentation, 1254
indoxacarb, 863
insecticidal bait, 863
insecticide resistance, 1548
toxicity, 863
Blissus occiduus,
Buchloë dactyloides, 1942
plant resistance, 1942
boll weevil, 311, 328, 348, 680
Anthonomus grandis, 348, 680
Anthonomus grandis grandis, 328
bait, 311
colonization, 328
cotton, 328, 680
kill strips, 348
malathion, 311
pheromone traps, 348
toxicity, 311
trapping, 680
boll weevil eradication,
Lygus lineolaris, 902
malathion, 902
tarnished plant bug, 902
Bollgard,
Bollgard II, 699
economic injury, 699
transgenics, 699
Bollgard II,
Bollgard, 699
economic injury, 699
transgenics, 699
bollworm,
Bacillus thuringiensis, 137
Bt resistance, 137
gene frequency, 137
boric acid,
cyfluthrin, 1362
German cockroach, 1362
integrated pest management (IPM), 1362
swine production, 1362
botanical insecticides,
Cryptolestes, 1005
peas, 1005
Sitophilus, *1005*
traps, 1005
Tribolium, 1005
botanicals,
feeding-deterrence, 1594
Rhyzopertha dominica, 1594
Sitophilus oryzae, 1594
toxicity, 1594
Tribolium castaneum, 1594
Brahman breeding,
Hematobia irritans irritans, 892
IGR-bolus, 892
innate fly resistance, 892
mechanical trap, 892
pasture rotation, 892
Brassica oleracea,
host plant resistance, 143
Mamestra brassicae, 143
Pieris brassicae, 143
Pieris rapae, 143
bruchid resistance,
Callosobruchus spp., 207
Ceratotropis, 207
rice bean, 207
Vigna spp., 207
Bt corn,
host plant resistance, 935
Ostrinia furnacalis, 935
transgenic plants, 935
Bt crops,
Bacillus thuringiensis, 1031
genetically modified crops, 1031
refuge, 1031
resistance, 1031
Bt resistance,
Bacillus thuringiensis, 137
bollworm, 137
gene frequency, 137
Buchloë dactyloides,
Blissus occiduus, 1942
plant resistance, 1942
budburst phenology,
bioassay, 377
Choristoneura occidentalis, 377
defoliation, 377
Pseudotsuga menziesii, 377
resistance mechanisms, 377

C

C. formosanus,
foraging, 88
fractal analysis, 88
R. flavipes, 88
tunnel geometry, 88
callaloo,
Amaranthus spp., 798
Herpetogramma bipunctalis, 798
Spodoptera spp., 798
Spoladea recurvalis, 798
Callosobruchus maculatus,
cold storage, 1952
cowpea weevil, 1952
dried beans, 1952
postharvest disinfestation, 1952
Callosobruchus spp.,
bruchid resistance, 207
Ceratotropis, 207
rice bean, 207
Vigna spp., 207
Calomycterus setarius,
Cerotoma trifurcata, 1168
photosynthesis, 1168
soybean, 1168
cannibalism,
Helicoverpa zea, 931
resistance management, 931
transgenic corn, 931
canopy structure,
Gossypium hirsutum, 730
Helicoverpa, 730
square loss, 730
terminal removal, 730
yield, 730
Carabidae,
corn, 1805
imidacloprid, 1805
Orius, 1805
predator, 1805
carabids,
Ataenius spretulus, 792
golf course, 792
milky disease, 792
Paenibacillus sp., 792
staphylinids, 792
carbaryl,
food-based bait, 584
grasshopper, 584
integrated pest management (IPM), 584
oil, 584
carotenoid,
chlorophyll, 475
Diuraphis noxia, 475
plant resistance, 475
Russian wheat aphid, 475
wheat, 475
cassava, 1905, 1915
cyanogenic potential, 1905, 1915
Cyrtomenus bergi, 1905, 1915
plant resistance, 1905, 1915
secondary plant metabolite, 1905, 1915
catnip,
nepetalactone, 1275
Reticulitermes, 1275
soil barrier, 1275
termite, 1275

Ceratitis capitata, 300, 615, 662, 1108, 1132, 1405, 1711
Anastrepha fraterculus, 1108
Argentina, 1108
bioinvasion, 1711
cold treatment, 300
dominant lethal mutation, 615
GF-120, 1405
host plants, 1108
mating competitiveness, 1132
Mediterranean fruit fly, 662
metamorphosis, 662
microsatellites, 1711
model, 300
mtDNA, 1711
phloxine B, 662
photoinsecticides, 662
polymerase chain reaction-restriction fragment length polymorphism, 1711
quality control, 615
sterile insect technique, 615, 1132, 1405
Tephritidae, 1108
tsl, 615
Ceratotropis,
bruchid resistance, 207
Callosobruchus spp., 207
rice bean, 207
Vigna spp., 207
cereal aphids, 975, 1585, 1792
conservation tillage, 1792
conventional tillage, 1792
Lysiphlebus testaceipes, 975
natural enemy threshold, 975
Rhopalosiphum padi, 1585, 1792
Schizaphis graminum, 1585
sequential sampling, 975, 1585
winter wheat, 1585
Cerotoma trifurcata,
Calomycterus setarius, 1168
photosynthesis, 1168
soybean, 1168
CheckMite+,
Apis mellifera, 259
Apistan, 259
honey bee, 259
tebufenpyrad, 259
varroa, 259
chemical control,
Delia antiqua, 1494
insect growth regulator, 1494
insect pest management, 1494
Chilo suppressalis,
Bacillus thuringiensis, 1838
China, 1838
susceptibility, 1838
China, 1322, 1838
Bacillus thuringiensis, 1322, 1838
Chilo suppressalis, 1838
Helicoverpa armigera, 1322
susceptibility, 1838
transgenic cotton, 1322
chitin synthesis inhibitor,
Australia, 1828
bait technology, 1828
chlorfluazuron, 1828
subterranean termites, 1828
chitinase *SH1*,
alginate beads, 555
bacteriophages, 555
Enterobacter cloacae, 555
Epilachna vigintioctopunctata, 555
chlorfluazuron,
Australia, 1828
bait technology, 1828
chitin synthesis inhibitor, 1828
subterranean termites, 1828
chlorophyll,
carotenoid, 475
Diuraphis noxia, 475
plant resistance, 475
Russian wheat aphid, 475
wheat, 475
chlorpyrifos,
acephate, 645
lambda-cyhalothrin, 645
methomyl, 645
Solenopsis invicta, 645
toxicology, 645
chlorpyrifos-methyl,
beetles, 525
psocids, 525
sorghum, 525
synergized bifenthrin, 525
choice tests,
Host plant resistance, 53
no-choice tests, 53
Odontota dorsalis, 53
Robinia pseudoacacia L., 53
Choristoneura fumiferana,
Bacillus thuringiensis, 280
Double application, 280
spruce budworm, 280
sublethal effects, 280
Choristoneura occidentalis, 377, 783
bioassay, 377
budburst phenology, 377
defoliation, 377
fertilization, 783
foliar nutrients, 783
Laccaria bicolor, 783
larval feeding preferences, 783
minerals, 783
Pseudotsuga menziesii, 377
resistance mechanisms, 377
Choristoneura rosaceana,
Bacillus thuringiensis, 879
insecticide resistance, 879
orchard pest management, 879
Pandemis limitata, 879
spinosad, 879
chronic,
Diatraea grandiosella, 12
microsporidia, 12
sublethal, 12
vertical transmission, 12
Chronic infection,
disease, 272
SPF, 272
technique, 272
Chrysomelidae, 609, 1738
Coleoptera, 609
Cry3Bb1, 1738
degree-days, 609
Diabrotica virgifera virgifera, 1738
insect resistance management, 1738
temperature, 609
western corn rootworm, 1738
Cirrospilus,
citrus, 577
dynamics, 577
parasitism, 577
peelminer, 577
citrullus lanatus,
Anasa tristis, 64
Cucurbitaceae, 64
Hemiptera, 64
citrus, 577, 811, 1693
Anastrepha fraterculus, 1693
antibiosis, 811
antifeedant, 811
Cirrospilus, 577
cover crops, 811
Diaprepes abbreviatus, 811
dynamics, 577
host status, 1693
oviposition behavior, 1693
parasitism, 577
peelminer, 577
pest status, 1693
citrus pests,
fruit fly lures, 1732
McPhail Trap, 1732
nontarget insects, 1732
Tephritidae, 1732
Citrus sinensis,
augmentative biological control, 1375
heteronomous hyperparasitoid, 1375
Citrus thrips,
molasses bait, 629
phloxine B, 629
photoactive dye, 629
Scirtothrips citri, 629
Cnephasia jactatana,
age, 592
mating delay, 592

reproductive potential, 592
Tortricidae, 592
cocoon penetration,
Hyposoter didymator, 1054
IGRs, 1054
naturalyte, 1054
parasitoid, 1054
pupae, 1054
Coffee tree,
economic damage threshold, 1473
estimation of losses, 1473
Hypothenemus hampei, 1473
Togo, 1473
cold storage,
Callosobruchus maculatus, 1952
cowpea weevil, 1952
dried beans, 1952
postharvest disinfestation, 1952
cold treatment,
Ceratitis capitata, 300
model, 300
Coleoptera, 609, 1126, 1704, 1770
aggregation pheromone, 1126
Chrysomelidae, 609
Curculionidae, 1126
degree-days, 609
remote sensing, 1770
Scarabaeidae, 1704, 1770
Scyphophorus acupunctatus, 1126
soil insects, 1704
sound, 1704, 1770
temperature, 609
turfgrass, 1704, 1770
colonization,
Anthonomus grandis grandis, 328
boll weevil, 328
cotton, 328
colony management,
Apis mellifera, 547
greenhouse, 547
honey bee, 547
pollination, 547
tomatoes, 547
common sugar,
Bacillus thuringiensis subsp. *israelensis*, 1039
Bengal gram, 1039
Luria Bertani, 1039
mosquitoes, 1039
potato, 1039
community ecology,
pet food, 1958
stored-product insects, 1958
trapping, 1958
compensation,
cotton, 1174
Helicoverpa zea, 1174
simulation, 1174
tolerance, 1174
computer simulation, 156, 173
Bacillus thuringiensis, 156
Bacillus thuringiensis, 173
resistance management, 156, 173
transgenic crops, 156, 173
concentration.,
Bactrocera cucurbitae, 231
food attractants, 231
protein hydrolysates, 231
Conotrachelus nenuphar, 826, 1399
attractants, 826
disinfestation, 1399
gamma radiation, 1399
monitoring, 826
odor, 826
phytosanitary treatment, 1399
trap trees, 826
conservation biological control,
host location, 1412
imidacloprid, 1412
Popillia japonica, 1412
sublethal effects, 1412
conservation tillage,
cereal aphids, 1792
conventional tillage, 1792
Rhopalosiphum padi, 1792
contact toxicity, 990, 1967
development inhibition, 990
d-Limonene, 990
feeding deterrent, 990
fumigant toxicity, 990
ovicide, 990
spinosad, 1967
stored-product insects, 1967
surface treatments, 1967
contour mapping,
IPM, 1617
semivariogram, 1617
spatial analysis, 1617
Trogoderma variabile, 1617
control, 126, 510, 844
Alphitobius diaperinus, 126
Argentine ant, 844
diatomaceous earth, 510
field trial, 126
foraging range, 844
insect growth regulator, 126
Linepithema humile, 844
maize weevil, 510
poultry house, 126
pyrethroid, 126
resurgence, 844
rice weevil, 510
Sitophilus, 510
stored grain, 510
Controlled atmosphere,
lettuce, 1100
postharvest pest control, 1100
postharvest quality, 1100
vacuum, 1100
conventional tillage,
cereal aphids, 1792
conservation tillage, 1792
Rhopalosiphum padi, 1792
Coptotermes formosanus, 106, 117, 456, 1526
deltamethrin, 1526
disodium octaborate tetrahydrate, 1526
exposure duration, 456
foam, 117
imidacloprid, 117
incipient colonies, 106
nonrepellent termiticides, 456
nutrition, 106
reproduction, 106
Reticulitermes flavipes, 1526
termite, 117
trees, 117
wood species, 106
Coptotermes formosanus Shiraki,
fipronil, 461
horizontal transmission, 461
repellency, 461
toxicity, 461
corn,
Carabidae, 1805
imidacloprid, 1805
Orius, 1805
predator, 1805
corn earworm,
cotton bollworm, 909
cross-resistance, 909
Helicoverpa, 909
host plant resistance, 909
Zea mays, 909
corn rootworm, 441, 669, 714, 1153
adult emergence, 714
bait, 441
cucurbitacin E-glycoside, 441
Diabrotica, 669
Diabrotica barberi, 1153
root injury, 714
soil insecticide rates, 714
Diabrotica undecimpunctata howardi, 441
Diabrotica virgifera virgifera, 1153
insect control, 441
larval instar determination, 1153
larval sampling, 1153
mitochondrial DNA, 669
species diagnostics, 669
cost analysis,
augmentative biological control, 1675
integrated pest management, 1675
natural enemies, 1675
ornamentals, 1675

costs,
Lymantria dispar L., 1448
spread, 1448
cotton, 328, 468, 680, 748, 805, 1174
aldicarb, 748
Anthonomus grandis, 680
Anthonomus grandis grandis, 328
boll weevil, 328, 680
colonization, 328
compensation, 1174
Frankliniella fusca, 748
Helicoverpa armigera, 468
Helicoverpa zea, 1174
insecticide mixtures, 468
insecticide resistance, 468
integrated pest management (IPM), 805
Kaolin, 805
particle films, 805
Pectinophora gossypiella, 805
pink bollworm, 805
precision placement, 748
simulation, 1174
Temik, 748
tobacco thrips, 748
tolerance, 1174
trapping, 680
West Africa, 468
cotton bollworm, 909, 1283
corn earworm, 909
cross-resistance, 909
cytochrome P450 monooxygenase, 1283
Helicoverpa, 909
Helicoverpa armigera, 1283
host plant resistance, 909
induction, 1283
tissue difference, 1283
Zea mays, 909
cotton integrated pest management,
lady beetles, 239
seasonality, 239
sticky trap, 239
trapping, 239
cottony cushion scale,
IGR, 1388
neonicotinoid, 1388
organophosphate, 1388
pyrethroid, 1388
cover crop, 811, 1426
antibiosis, 811
antifeedant, 811
Beta vulgaris, 1426
Citrus, 811
Diaprepes abbreviatus, 811
integrated control, 1426
soil insecticide, 1426
Tetanops myopaeformis, 1426
cowpea,
intraspecific variation, 489
pest management, 489
resistance, 489
seed beetle, 489
cowpea weevil,
Callosobruchus maculatus, 1952
cold storage, 1952
dried beans, 1952
postharvest disinfestation, 1952
cranberry,
Anthonomus musculus, 407
cranberry weevil, 407
plant compensation, 407
Vaccinium, 407
yield, 407
cranberry weevil,
Anthonomus musculus, 407
cranberry, 407
plant compensation, 407
Vaccinium, 407
yield, 407
crop phenology,
baculovirus, 649
maize, 649
natural enemies, 649
synthetic pesticides, 649
crop rotation, 737, 1872
Diabrotica virgifera virgifera, 737, 1872
insect resistance management, 1872
modeling, 1872
western corn rootworm, 737
cross-resistance, 909, 1548
Blattella germanica, 1548
corn earworm, 909
cotton bollworm, 909
cyclodienes, 1548
Fipronil, 1548
Helicoverpa, 909
host plant resistance, 909
insecticide resistance, 1548
Zea mays, 909
Cry1Ac,
Cry2Ab, 755
lepidopterous pests, 755
transgenic *Bacillus thuringiensis* cotton, 755
Cry2Ab,
Cry1Ac, 755
lepidopterous pests, 755
transgenic *Bacillus thuringiensis* cotton, 755
Cry3Bb1,
Chrysomelidae, 1738
Diabrotica virgifera virgifera, 1738
insect resistance management, 1738
western corn rootworm, 1738
Cryptolestes,
botanical insecticides, 1005
peas, 1005
Sitophilus, 1005
traps, 1005
Tribolium, 1005
Cryptophlebia leucotreta,
fecundity, 1724
fertility, 1724
gamma radiation, 1724
sterile insect technique, 1724
Cucurbitaceae,
Anasa tristis, 64
citrullus lanatus, 64
Hemiptera, 64
cucurbitacin E-glycoside,
bait, 441
corn rootworm, 441
Diabrotica undecimpunctata howardi, 441
insect control, 441
cue-lure,
melon fly, 1780
methyl eugenol, 1780
oriental fruit fly, 1780
spinosad, 1780
Culex,
Aedes, 1786
Bacillus thuringiensis variety *israelensis*, 1786
biocontrol, 1786
Ochlerotatus, 1786
Culex nigripalpus,
Aedes albopictus, 1559
Anopheles quadrimaculatus, 1559
toxicity, 1559
cultivar preference,
Fiji disease virus, 148
Perkinsiella saccharicida, 148
planthopper vector, 148
resistance, 148
sugar cane, 148
cultural control, 706, 850
aromatic cedar mulch, 850
biological control, 706
field study, 850
Linepithema humile, 850
Myiopharus macellus, 706
nesting sites, 850
sunflower beetle, 706
Zygogramma exclamationis, 706
Curculionidae,
aggregation pheromone, 1126
Coleoptera, 1126
Scyphophorus acupunctatus, 1126
cyanogenic potential, 1905, 1915
cassava, 1905, 1915
Cyrtomenus bergi, 1905, 1915
plant resistance, 1905, 1915

secondary plant metabolite, 1905, 1915
cyclodienes,
Blattella germanica, 1548
cross-resistance, 1548
Fipronil, 1548
insecticide resistance, 1548
Cydia pomonella,
methoxyfenozide, 623
orientation tube, 623
sex pheromone, 623
wind tunnel, 623
cyfluthrin,
boric acid, 1362
German cockroach, 1362
integrated pest management (IPM), 1362
swine production, 1362
λ-cyhalothrin,
indoxacarb, 1230
Plutella xylostella, 1230
reduced-risk insecticide, 1230
Trichoplusia ni, 1230
cyromazine,
diflubenzuron, 1300
insect growth regulator (IGR), 1300
methoprene, 1300
selection, 1300
Cyrtomenus bergi, 1905, 1915
cassava, 1905, 1915
cyanogenic potential, 1905, 1915
plant resistance, 1905, 1915
secondary plant metabolite, 1905, 1915
Cyrtomenus ciliatus,
Arachis hypogaea, 1142
economic injury, 1142
Pangaeus bilineatus, 1142
Sehirus cinctus, 1142
cytochrome P450 monooxygenase,
cotton bollworm, 1283
Helicoverpa armigera, 1283
induction, 1283
tissue difference, 1283

D

Dasineura oxycoccana,
monitoring, 1821
traps, 1821
Vaccinium spp., 1821
defoliation,
bioassay, 377
budburst phenology, 377
Choristoneura occidentalis, 377
Pseudotsuga menziesii, 377
resistance mechanisms, 377
degree-days,
Chrysomelidae, 609
Coleoptera, 609
temperature, 609
Delia antiqua,
chemical control, 1494
insect growth regulator, 1494
insect pest management, 1494
deltamethrin,
Coptotermes formosanus, 1526
disodium octaborate tetrahydrate, 1526
Reticulitermes flavipes, 1526
Dendroctonus rufipennis,
AK (south-central), 31
antiaggregation, 31
Bark beetles, 31
Picea sp., 31
semiochemicals, 31
Dermestes maculatus,
Alphitobius diaperinus, 1602
Beauveria bassiana, 1602
poultry, 1602
development, 43, 1119
host plant suitability, 43
Lymantria monacha, 43
Nysius huttoni, 1119
reproduction, 1119
survival, 43, 1119
temperature, 1119
development inhibition,
contact toxicity, 990
d-Limonene, 990
feeding deterrent, 990
fumigant toxicity, 990
ovicide, 990
Diabrotica, 131, 669
corn rootworm, 669
esterase, 131
inheritance, 131
mitochondrial DNA, 669
reciprocal cross, 131
species diagnostics, 669
western corn rootworm, 131
Diabrotica barberi,
corn rootworm, 1153
Diabrotica virgifera virgifera, 1153
larval instar determination, 1153
larval sampling, 1153
Diabrotica barberi Smith Lawrence,
Diabrotica virgifera virgifera LeConte, 1420
northern corn rootworm, 1420
sampling efficiency, 1420
western corn rootworm, 1420
Diabrotica undecimpunctata howardi,
bait, 441
corn rootworm, 441
cucurbitacin E-glycoside, 441
insect control, 441
Diabrotica virgifera,
mark-recapture, 1750
rubidium, 1750
western corn rootworm, 1750
Zea mays, 1750
Diabrotica virgifera virgifera, 599, 737, 1153, 1738, 1855, 1872
Chrysomelidae, 1738
corn rootworm, 1153
crop rotation, 737, 1872
Cry3Bb1, 1738
Diabrotica barberi, 1153
esterase, 1855
insect resistance management, 1738, 1872
insecticide resistance, 1855
larval instar determination, 1153
larval movement, 599
larval sampling, 1153
methyl parathion, 1855
modeling, 1872
organophosphate, 1855
resistance management, 599
western corn rootworm, 599, 737, 1738
Diabrotica virgifera virgifera LeConte, 1420, 1530
Diabrotica barberi Smith Lawrence, 1420
insecticidal traits, 1530
northern corn rootworm, 1420
population dynamics, 1530
resistance management, 1530
sampling efficiency, 1420
western corn rootworm, 1420
diapause,
migration, 1433
pest management, 1433
phenology, 1433
resistance management, 1433
Diaprepes abbreviatus,
antibiosis, 811
antifeedant, 811
Citrus, 811
cover crops, 811
diatomaceous earth,
control, 510
maize weevil, 510
rice weevil, 510
Sitophilus, 510
stored grain, 510
Diatraea grandiosella,
chronic, 12
microsporidia, 12
sublethal, 12
vertical transmission, 12
diflubenzuron, 396, 1300
Bacillus thurengiensis var. *kurstaki*, 396
cyromazine, 1300

insect growth regulator (IGR), 1300
methoprene, 1300
Orgyia pseudotsugata, 396
selection, 1300
tebufenozide, 396
disease,
Chronic infection, 272
SPF, 272
technique, 272
disinfestation,
Conotrachelus nenuphar, 1399
gamma radiation, 1399
phytosanitary treatment, 1399
disodium octaborate tetrahydrate,
Coptotermes formosanus, 1526
deltamethrin, 1526
Reticulitermes flavipes, 1526
Diuraphis noxia, 194, 214, 352, 475, 673
carotenoid, 475
chlorophyll, 475
host plant resistance, 352, 673
light modulation, 194
mechanisms of resistance, 214
photosynthesis, 194
plant physiology, 194
plant resistance, 475
Russian wheat aphid, 194, 475, 673
wheat, 194, 214, 352, 475, 673
d-Limonene,
contact toxicity, 990
development inhibition, 990
feeding deterrent, 990
fumigant toxicity, 990
ovicide, 990
dog,
American cockroach, 1259
drywood termites, 1259
subterranean termites, 1259
termite detection, 1259
termite-damaged wood, 1259
dominant lethal mutation,
Ceratitis capitata, 615
quality control, 615
sterile insect technique, 615
tsl, 615
dosage compensation,
arrhenotokous, 1626
haplodiploid resistance, 1626
resistance in natural enemies, 1626
sexual dimorphism resistance, 1626
Double application,
Bacillus thuringiensis, 280
Choristoneura fumiferana, 280
spruce budworm, 280
sublethal effects, 280
dried beans,
Callosobruchus maculatus, 1952
cold storage, 1952
cowpea weevil, 1952
postharvest disinfestation, 1952
dried fruits,
Heat treatments, 519
Indianmeal moth, 519
postharvest, 519
radio frequency, 519
tree nuts, 519
dry weight loss,
bioassay, 220
host plant resistance, 220
kernel damage, 220
Rhyzopertha dominica, 220
stored-products, 220
drywood termites,
American cockroach, 1259
dog, 1259
subterranean termites, 1259
termite detection, 1259
termite-damaged wood, 1259
dynamics,
Cirrospilus, 577
citrus, 577
parasitism, 577
peelminer, 577

E

Ebeling choice box,
Blattella germanica, 863
field performance, 863
indoxacarb, 863
insecticidal bait, 863
toxicity, 863
economic damage threshold,
Coffee tree, 1473
estimation of losses, 1473
Hypothenemus hampei, 1473
Togo, 1473
economic injury, 699, 1142
Arachis hypogaea, 1142
Bollgard, 699
Bollgard II, 699
Cyrtomenus ciliatus, 1142
Pangaeus bilineatus, 1142
Sehirus cinctus, 1142
transgenics, 699
economic threshold, 1160, 1500
EIL, 1160
Epilachna varivestis, 1160
integrated pest management, 1500
Phaseolus vulgaris, 1160
strawberry, 1500
tarnished plant bug, 1500
edge effect,
Blattella germanica, 1832
exploratory behavior, 1832
efficacy,
Bacillus thuringiensis, 388
experimental design, 388
replication, 388
spruce budworm, 388
efficiency,
Blattella germanica, 1254
fipronil gel bait, 1254
fragmentation, 1254
egg predators,
insecticide, 71
pests, 71
Zea mays, 71
egg sampling,
Lambdina fiscellaria, 768
light trap, 768
monitoring, 768
outbreak detection, 768
oviposition trap, 768
eggs,
maturation, 341
oogenesis, 341
sugar, 341
yeast hydrolysate, 341
EIL,
economic threshold, 1160
Epilachna varivestis, 1160
Phaseolus vulgaris, 1160
elm leaf beetle,
abamectin, 1455
imidacloprid, 1455
integrated pest management, 1455
Xanthogaleruca luteola, 1455
Empoasca fabae, 690, 1184
Anticarsia gemmatalis, 690
genetically modified soybeans, 690
Medicago sativa, 1184
Nezara viridula, 690
Roundup Ready soybeans, 690
Spissistilus festinus, 690
strip harvesting, 1184
Encarsia,
Bemisia tabaci, 543
Eretmocerus, 543
LED-plastic cup trap, 543
yellow sticky card trap, 543
Endopiza viteana,
leafhoppers, 1221
Maladera castanea, 1221
vineyards, 1221
Enterobacter cloacae,
alginate beads, 555
bacteriophages, 555
chitinase *SH1*, 555
Epilachna vigintioctopunctata, 555
entomopathogenic nematodes, 287, 1076
application methodology, 287
Heterorhabditis, 1076

Heterorhabditis bacteriophora, 287
insecticides, 1076
Maladera castanea, 1076
Steinernema, 1076
Steinernema spp., 287
Vitis, 287
Eoreuma loftini, 570, 950, 1929
areawide pest management, 1929
biological control, 570
fitness, 570
Galanthus nivalis agglutinin, 570, 950
Galanthus nivalis L., 950
host plant resistance, 950, 1929
life table parameters, 950
nontarget effects, 570
sugarcane, 1929
Epilachna varivestis,
economic threshold, 1160
EIL, 1160
Phaseolus vulgaris, 1160
Epilachna vigintioctopunctata,
alginate beads, 555
bacteriophages, 555
chitinase *SH1*, 555
Enterobacter cloacae, 555
Eretmocerus,
Bemisia tabaci, 543
Encarsia, 543
LED-plastic cup trap, 543
yellow sticky card trap, 543
esfenvalerate,
insect genetics, 1021
insecticide resistance, 1021
LdVssc1, 1021
Para, 1021
essential oil,
mode of action, 1479
natural fumigant, 1479
natural insecticide, 1479
Trialeurodes vaporariorum, 1479
esterase, 131, 322, 1855
Bemisia tabaci, 322
biotype, 322
Diabrotica, 131
Diabrotica virgifera virgifera, 1855
inheritance, 131
insecticide resistance, 1855
methyl parathion, 1855
organophosphate, 1855
reciprocal cross, 131
virus, 322
western corn rootworm, 131
estimation of losses,
Coffee tree, 1473
economic damage threshold, 1473
Hypothenemus hampei, 1473
Togo, 1473
European corn borer, 361, 503, 957
Bacillus thuringiensis corn, 957
imidacloprid, 503
Indianmeal moth, 503
millet, 361
Ostrinia nubilalis, 361
permethrin, 957
Plodia interpunctella, 503
refuge, 361
South Dakota, 957
thiamethoxam, 503
trap crop, 361
univoltine ecotype, 957
yield, 957
experimental design,
Bacillus thuringiensis, 388
efficacy, 388
replication, 388
spruce budworm, 388
exploratory behavior,
Blattella germanica, 1832
edge effect, 1832
exposure duration,
Coptotermes formosanus, 456
nonrepellent termiticides, 456
extra binomial variation,
bioassay, 983
generalized linear mixed model, 983
probit analysis, 983
spray oils, 983

F

fecundity,
Cryptophlebia leucotreta, 1724
fertility, 1724
gamma radiation, 1724
sterile insect technique, 1724
feeding deterrent,
contact toxicity, 990
development inhibition, 990
d-Limonene, 990
fumigant toxicity, 990
ovicide, 990
feeding-deterrence,
botanicals, 1594
Rhyzopertha dominica, 1594
Sitophilus oryzae, 1594
toxicity, 1594
Tribolium castaneum, 1594
fertility,
Cryptophlebia leucotreta, 1724
fecundity, 1724
gamma radiation, 1724
sterile insect technique, 1724
fertilization,
Choristoneura occidentalis, 783
foliar nutrients, 783
Laccaria bicolor, 783
larval feeding preferences, 783
minerals, 783
Festuca arundinacea,
Lolium perenne, 1898
Poa pratensis, 1898
Rhizotrogus majalis, 1898
tolerance, 1898
field insecticide tests,
laboratory bioassays, 1463
Rhagoletis pomonella, 1463
field performance,
Blattella germanica, 863
Ebeling choice box, 863
indoxacarb, 863
insecticidal bait, 863
toxicity, 863
field study,
aromatic cedar mulch, 850
cultural control, 850
Linepithema humile, 850
nesting sites, 850
field trial,
Alphitobius diaperinus, 126
control, 126
insect growth regulator, 126
poultry house, 126
pyrethroid, 126
Fiji disease virus,
cultivar preference, 148
Perkinsiella saccharicida, 148
planthopper vector, 148
resistance, 148
sugar cane, 148
fipronil, 461, 1548
Blattella germanica, 1548
Coptotermes formosanus Shiraki, 461
cross-resistance, 1548
cyclodienes, 1548
horizontal transmission, 461
insecticide resistance, 1548
repellency, 461
toxicity, 461
fipronil gel bait,
Blattella germanica, 1254
efficiency, 1254
fragmentation, 1254
fitness, 570, 1290
Bacillus thuringiensis, 1290
biological control, 570
Eoreuma loftini, 570
Galanthus nivalis agglutinin, 570
Helicoverpa armigera, 1290
nontarget effects, 570
resistance, 1290
transgenic cotton, 1290
fluorescence,
Aphis glycines, 188
gas exchange, 188
insect-plant relations, 188
photosynthesis, 188

foam,
Coptotermes formosanus, 117
imidacloprid, 117
termite, 117
trees, 117
foliar nutrients,
Choristoneura occidentalis, 783
fertilization, 783
Laccaria bicolor, 783
larval feeding preferences, 783
minerals, 783
food attractants,
Bactrocera cucurbitae, 231
concentration., 231
protein hydrolysates, 231
food-based bait,
carbaryl, 584
grasshopper, 584
integrated pest management (IPM), 584
oil, 584
foraging, 88, 871
Argentine ant, 871
C. formosanus, 88
fractal analysis, 88
granular bait, 871
Linepithema humile, 871
R. flavipes, 88
tunnel geometry, 88
foraging range,
Argentine ant, 844
control, 844
Linepithema humile, 844
resurgence, 844
formulation,
B. thuringiensis, 292
biopesticides, 292
spray drying, 292
suspensibility, 292
wettability, 292
fractal analysis,
C. formosanus, 88
foraging, 88
R. flavipes, 88
tunnel geometry, 88
fragmentation,
Blattella germanica, 1254
efficiency, 1254
fipronil gel bait, 1254
Frankliniella,
Geocoris punctipes, 1201
Orius insidiosus, 1201
spinosad, 1201
UV reflective mulch, 1201
Frankliniella fusca,
aldicarb, 748
Cotton, 748
precision placement, 748
Temik, 748
tobacco thrips, 748
Frankliniella occidentalis,
Aphis gossypii, 246
organosilicones, 246
Platynota stultana, 246
Pseudococcus maritimus, 246
Tetranychus pacificus, 246
thrips, 1
tomato spotted wilt virus, 1
Fraser fir,
aesthetic injury level, 1760
balsam fir, 1760
host plant phenology, 1760
integrated pest management, 1760
fruit fly lures,
citrus pests, 1732
McPhail Trap, 1732
nontarget insects, 1732
Tephritidae, 1732
fumigant toxicity,
contact toxicity, 990
development inhibition, 990
d-Limonene, 990
feeding deterrent, 990
ovicide, 990
funnel trap,
larval monitoring, 835
root weevil, 835

G

Galanthus nivalis agglutinin, 570, 950
biological control, 570
Eoreuma loftini, 570, 950
fitness, 570
Galanthus nivalis L., 950
host plant resistance, 950
life table parameters, 950
nontarget effects, 570
Galanthus nivalis L.,
Eoreuma loftini, 950
Galanthus nivalis agglutinin, 950
host plant resistance, 950
life table parameters, 950
gall,
Hemadas nubilipennis, 1245
lowbush blueberry, 1245
Vaccinium angustifolium, 1245
gamma radiation, 1399, 1724
Conotrachelus nenuphar, 1399
Cryptophlebia leucotreta, 1724
disinfestation, 1399
fecundity, 1724
fertility, 1724
phytosanitary treatment, 1399
sterile insect technique, 1724
garden symphylan,
bait sampling, 969
Scutigerella immaculata, 969
soil ecology, 969
soil sampling, 969
symphylid, 969
gas exchange,
Aphis glycines, 188
fluorescence, 188
insect-plant relations, 188
photosynthesis, 188
gene frequency,
Bacillus thuringiensis, 137
bollworm, 137
Bt resistance, 137
generalized linear mixed model,
bioassay, 983
extra binomial variation, 983
probit analysis, 983
spray oils, 983
generic Taylor coefficients,
ivy geranium, 482
presence-absence sampling, 482
twospotted spider mite, 482
genetic interaction,
resistance screening, 202
Schizaphis graminum, 202
synthetic hexaploid wheat, 202
genetically modified crops,
Bacillus thuringiensis, 1031
Bt crops, 1031
refuge, 1031
resistance, 1031
genetically modified soybeans,
Anticarsia gemmatalis, 690
Empoasca fabae, 690
Nezara viridula, 690
Roundup Ready soybeans, 690
Spissistilus festinus, 690
Geocoris punctipes,
Frankliniella, 1201
Orius insidiosus, 1201
spinosad, 1201
UV reflective mulch, 1201
German cockroach, 885, 1362
boric acid, 1362
cyfluthrin, 1362
integrated pest management (IPM), 1362
kdr-type resistance, 885
knockdown resistance, 885
multiplex polymerase chain reaction, 885
para, 885
swine production, 1362
GF-120,
Ceratitis capitata, 1405
sterile insect technique, 1405
GF-120 fruit fly bait,
Bactrocera cucurbitae, 1485
bait sprays, 1485
spinosad, 1485
glassy-winged sharpshooter, 264, 1849
Homalodisca coagulata, 1849
Homalodisca lacerta, 1849

insecticide resistance, 1849
Pierce's disease, 264
smoke-tree sharpshooter, 1849
vector, 264
xylem, 264
golf course,
Ataenius spretulus, 792
carabids, 792
milky disease, 792
Paenibacillus sp., 792
staphylinids, 792
Gossypium hirsutum,
canopy structure, 730
Helicoverpa, 730
square loss, 730
terminal removal, 730
yield, 730
grain elevator,
natural enemies, 996
population dynamics, 996
granular bait,
Argentine ant, 871
foraging, 871
Linepithema humile, 871
granular contact insecticide,
ant exclusion, 98
perimeter treatment, 98
repellency, 98
Grapholita molesta,
apple, 58
mating disruption, 58
microencapsulation, 58
sprayable pheromone, 58
grasshopper,
carbaryl, 584
food-based bait, 584
integrated pest management (IPM), 584
oil, 584
grasshopper density,
above- below-ground growth, 1193
Melanoplus sanguinipes, 1193
oilseed crop, 1193
small-grains, 1193
green apple aphid,
Aphis pomi, 1864
imidacloprid, 1864
mortality, 1864
toxicity, 1864
greenhouse,
Apis mellifera, 547
colony management, 547
honey bee, 547
pollination, 547
tomatoes, 547
groundnut,
Arachis spp., 1886
Helicoverpa, 1886
resistance, 1886
Spodoptera, 1886
Gypchek,
gypsy moth, 1662
phenolic glycosides, 1662
Populus, 1662
salicin, 1662
gypsy moth,
Gypchek, 1662
phenolic glycosides, 1662
Populus, 1662
salicin, 1662

H

haplodiploid resistance,
arrhenotokous, 1626
dosage compensation, 1626
resistance in natural enemies, 1626
sexual dimorphism resistance, 1626
heat tolerance,
heat treatment, 1345
methyl bromide alternative, 1345
Tribolium castaneum, 1345
heat treatment, 519, 1345
dried fruits, 519
heat tolerance, 1345
Indianmeal moth, 519
methyl bromide alternative, 1345
postharvest, 519
radio frequency, 519
tree nuts, 519
Tribolium castaneum, 1345
heifers,
beef cattle, 1612
Hematobia irritans, 1612
reproduction, 1612
weight gain, 1612
Helicoverpa, 730, 909, 1886
Arachis spp., 1886
canopy structure, 730
corn earworm, 909
cotton bollworm, 909
cross-resistance, 909
Gossypium hirsutum, 730
groundnut, 1886
host plant resistance, 909
resistance, 1886
Spodoptera, 1886
square loss, 730
terminal removal, 730
yield, 730
Zea mays, 909
Helicoverpa armigera, 468, 1283, 1290, 1322
Bacillus thuringiensis, 1290, 1322
China, 1322
cotton, 468
cotton bollworm, 1283
cytochrome P450 monooxygenase, 1283
fitness, 1290
induction, 1283
insecticide mixtures, 468
insecticide resistance, 468
resistance, 1290
tissue difference, 1283
transgenic cotton, 1290, 1322
West Africa, 468
Helicoverpa zea, 914, 925, 931, 1174, 1653
adult emergence, 925
Bacillus thuringiensis, 1653
Beauveria bassiana, 1653
cannibalism, 931
compensation, 1174
cotton, 1174
moderate dose, 914
Ostrinia nubilalis, 1653
plant resistance, 914
resistance management, 925, 931
simulation, 1174
sweet corn, 1653
tolerance, 1174
transgenic corn, 914, 925, 931
Hemadas nubilipennis,
gall, 1245
lowbush blueberry, 1245
Vaccinium angustifolium, 1245
Hematobia irritans, 1608, 1612
beef cattle, 1612
heifers, 1612
ivermectin, 1608
reproduction, 1612
sustained release bolus, 1608
weight gain, 1612
Hematobia irritans irritans,
Brahman breeding, 892
IGR-bolus, 892
innate fly resistance, 892
mechanical trap, 892
pasture rotation, 892
Hemiptera, 64, 251
Anasa tristis, 64
ants, 251
biological control, 251
citrullus lanatus, 64
Cucurbitaceae, 64
insecticides, 251
territoriality, 251
hemlock,
Adelges tsugae, 763
biological control, 763
hemlock woolly adelgid, 763
Pseudoscymnus tsugae, 763
Scymnus ningshanensis, 763
hemlock woolly adelgid,
Adelges tsugae, 763
biological control, 763
hemlock, 763

Pseudoscymnus tsugae, 763
Scymnus ningshanensis, 763
Herpetogramma bipunctalis,
Amaranthus spp., 798
callaloo, 798
Spodoptera spp., 798
Spoladea recurvalis, 798
heterobeltiosis,
antibiosis, 1935
heterosis, 1935
host plant resistance, 1935
hybrid rice, 1935
heteronomous hyperparasitoid,
augmentative biological control, 1375
Citrus sinensis, 1375
Heterorhabditis,
entomopathogenic nematodes, 1076
insecticides, 1076
Maladera castanea, 1076
Steinernema, 1076
Heterorhabditis bacteriophora,
application methodology, 287
entomopathogenic nematodes, 287
Steinernema spp., 287
Vitis, 287
heterosis,
antibiosis, 1935
heterobeltiosis, 1935
host plant resistance, 1935
hybrid rice, 1935
Homalodisca coagulata,
glassy-winged sharpshooter, 1849
Homalodisca lacerta, 1849
insecticide resistance, 1849
smoke-tree sharpshooter, 1849
Homalodisca lacerta,
glassy-winged sharpshooter, 1849
Homalodisca coagulata, 1849
insecticide resistance, 1849
smoke-tree sharpshooter, 1849
honey bee, 259, 547, 1510
Allium cepa, 1510
Apis mellifera, 259, 547
Apistan, 259
CheckMite+, 259
colony management, 547
greenhouse, 547
honey bee behavior, 1510
onion, 1510
pollination, 547, 1510
tebufenpyrad, 259
tomatoes, 547
varroa, 259
honey bee behavior,
Allium cepa, 1510
honey bees, 1510
onion, 1510
pollination, 1510
horizontal transmission,
Coptotermes formosanus Shiraki, 461
fipronil, 461
repellency, 461
toxicity, 461
host location, 777, 1412
conservation biological control, 1412
imidacloprid, 1412
introduced basswood thrips, 777
olfactory cues, 777
pear thrips, 777
pest monitoring, 777
Popillia japonica, 1412
sublethal effects, 1412
visual cues, 777
host plant,
Bemisia argentifolii, 1315
bifenthrin resistance, 1315
inheritance, 1315
host plant phenology,
aesthetic injury level, 1760
balsam fir, 1760
Fraser fir, 1760
integrated pest management, 1760
host plant resistance, 53, 143, 220, 352, 673, 909, 935, 950, 1577, 1929, 1935
antibiosis, 1577, 1935
antixenosis, 1577
areawide pest management, 1929
bioassay, 220
Brassica oleracea, 143
Bt corn, 935
choice tests, 53
corn earworm, 909
cotton bollworm, 909
cross-resistance, 909
Diuraphis noxia, 352, 673
dry weight loss, 220
Eoreuma loftini, 950, 1929
Galanthus nivalis agglutinin, 950
Galanthus nivalis L., 950
Helicoverpa, 909
heterobeltiosis, 1935
heterosis, 1935
hybrid rice, 1935
kernel damage, 220
life table parameters, 950
Mamestra brassicae, 143
no-choice tests, 53
Odontota dorsalis, 53
Ostrinia furnacalis, 935
Phaseolus vulgaris, 1577
Pieris brassicae, 143
Pieris rapae, 143
Rhyzopertha dominica, 220
Robinia pseudoacacia L., 53
Russian wheat aphid, 673
stored-products, 220
sugarcane, 1929
Thrips palmi, 1577
transgenic plants, 935
wheat, 352, 673
Zea mays, 909
host plant suitability,
development, 43
Lymantria monacha, 43
survival, 43
host plants,
Anastrepha fraterculus, 1108
Argentina, 1108
Ceratitis capitata, 1108
Tephritidae, 1108
host status,
Anastrepha fraterculus, 1693
citrus, 1693
oviposition behavior, 1693
pest status, 1693
house fly,
Alphitobius diaperinus, 1352
lesser mealworm, 1352
Musca domestica, 1352
poultry pest management, 1352
human urine,
Anastrepha, 334
attractants, 334
Tephritidae, 334
hybrid rice,
antibiosis, 1935
heterobeltiosis, 1935
heterosis, 1935
host plant resistance, 1935
hydrogen phosphide fumigation,
Mayetiola destructor (Say), 1340
pressure, 1340
Hymenoptera,
beneficial, 1083
integrated pest management, 1083
pesticide, 1083
Trichogramma, 1083
Hyposoter didymator,
cocoon penetration, 1054
IGRs, 1054
naturalyte, 1054
parasitoid, 1054
pupae, 1054
Hypothenemus hampei,
Coffee tree, 1473
economic damage threshold, 1473
estimation of losses, 1473
Togo, 1473

I

IGR,
cottony cushion scale, 1388
neonicotinoid, 1388
organophosphate, 1388
pyrethroid, 1388
IGR-bolus,
Brahman breeding, 892
Hematobia irritans irritans, 892
innate fly resistance, 892
mechanical trap, 892
pasture rotation, 892
IGRs,
cocoon penetration, 1054
Hyposoter didymator, 1054
naturalyte, 1054
parasitoid, 1054
pupae, 1054
imidacloprid, 117, 503, 1412, 1455, 1805, 1864
abamectin, 1455
Aphis pomi, 1864
Carabidae, 1805
conservation biological control, 1412
Coptotermes formosanus, 117
corn, 1805
elm leaf beetle, 1455
European corn borer, 503
foam, 117
green apple aphid, 1864
host location, 1412
Indianmeal moth, 503
integrated pest management, 1455
mortality, 1864
Orius, 1805
Plodia interpunctella, 503
Popillia japonica, 1412
predator, 1805
sublethal effects, 1412
termite, 117
thiamethoxam, 503
toxicity, 1864
trees, 117
Xanthogaleruca luteola, 1455
incipient colonies,
Coptotermes formosanus, 106
nutrition, 106
reproduction, 106
wood species, 106
Indianmeal moth, 503, 519
dried fruits, 519
European corn borer, 503
Heat treatments, 519
imidacloprid, 503
Plodia interpunctella, 503
postharvest, 519
radio frequency, 519
thiamethoxam, 503
tree nuts, 519
indoxacarb, 863, 1230
Blattella germanica, 863
λ-cyhalothrin, 1230
Ebeling choice box, 863
field performance, 863
insecticidal bait, 863
Plutella xylostella, 1230
reduced-risk insecticide, 1230
toxicity, 863
Trichoplusia ni, 1230
induction,
cotton bollworm, 1283
cytochrome P450 monooxygenase, 1283
Helicoverpa armigera, 1283
tissue difference, 1283
infestation,
Nilaparvata lugens, 1798
nutrient uptake, 1798
rice root, 1798
inheritance, 131, 1315
Bemisia argentifolii, 1315
bifenthrin resistance, 1315
Diabrotica, 131
esterase, 131
host plant, 1315
reciprocal cross, 131
western corn rootworm, 131
innate fly resistance,
Brahman breeding, 892
Hematobia irritans irritans, 892
IGR-bolus, 892
mechanical trap, 892
pasture rotation, 892
insect control,
bait, 441
corn rootworm, 441
cucurbitacin E-glycoside, 441
Diabrotica undecimpunctata howardi, 441
insect genetics,
esfenvalerate, 1021
insecticide resistance, 1021
LdVssc1, 1021
Para, 1021
insect growth regulator, 126, 1300, 1307, 1494
Alphitobius diaperinus, 126
Bemisia, 1307
chemical control, 1494
control, 126
cyromazine, 1300
Delia antiqua, 1494
diflubenzuron, 1300
field trial, 126
insect pest management, 1494
methoprene, 1300
poultry house, 126
pyrethroid, 126
pyriproxyfen, 1307
resistance, 1307
selection, 1300
whitefly, 1307
insect pest management,
chemical control, 1494
Delia antiqua, 1494
insect growth regulator, 1494
insect resistance management, 1738, 1872
Chrysomelidae, 1738
crop rotation, 1872
Cry3Bb1, 1738
Diabrotica virgifera virgifera, 1738, 1872
modeling, 1872
western corn rootworm, 1738
insecta, 370, 413
biological control, 370
nutrition, 370
Ostrinia nubilalis, 413
Phoracantha recurva, 370
Phoracantha semipunctata, 370
pollen, 370
sweet corn, 413
Zea mays, 413
insecticidal activity,
nucleopolyhedrovirus, 1668
optical brightener, 1668
Spodoptera exigua, 1668
synergism, 1668
insecticidal bait,
Blattella germanica, 863
Ebeling choice box, 863
field performance, 863
indoxacarb, 863
toxicity, 863
insecticidal traits,
Diabrotica virgifera virgifera LeConte, 1530
population dynamics, 1530
resistance management, 1530
insecticide, 71, 251, 856, 1076
ants, 251, 856
biological control, 251
egg predators, 71
entomopathogenic nematodes, 1076
hemiptera, 251
Heterorhabditis, 1076
Maladera castanea, 1076
pests, 71
Steinernema, 1076
Tapinoma melanocephalum, 856
territoriality, 251
toxic baits, 856
Zea mays, 71
insecticide mixtures,
cotton, 468
Helicoverpa armigera, 468
insecticide resistance, 468
West Africa, 468

insecticide resistance, 468, 875, 879, 1016, 1021, 1548, 1849, 1855
Aphis gossypii, 875
Bacillus thuringiensis, 879
bioassay, 1016
Blattella germanica, 1548
Choristoneura rosaceana, 879
cotton, 468
cross-resistance, 1548
cyclodienes, 1548
Diabrotica virgifera virgifera, 1855
esfenvalerate, 1021
esterase, 1855
Fipronil, 1548
glassy-winged sharpshooter, 1849
Helicoverpa armigera, 468
Homalodisca coagulata, 1849
Homalodisca lacerta, 1849
insect genetics, 1021
insecticide mixtures, 468
LdVssc1, 1021
methoxychlor, 1016
methyl parathion, 1855
Musca domestica, 1016
orchard pest management, 879
organophosphate, 1855
Pakistan, 875
Pandemis limitata, 879
Para, 1021
permethrin, 1016
pyrethroids, 875
smoke-tree sharpshooter, 1849
spinosad, 879
stirofos, 1016
West Africa, 468
insect-plant relations,
Aphis glycines, 188
fluorescence, 188
gas exchange, 188
photosynthesis, 188
integrated control,
Beta vulgaris, 1426
cover crop, 1426
soil insecticide, 1426
Tetanops myopaeformis, 1426
integrated pest management (IPM), 21, 401, 584, 805, 1083, 1362, 1455, 1500, 1647, 1675, 1760
abamectin, 1455
aesthetic injury level, 1760
Aethina tumida, 1647
Apis mellifera, 1647
augmentative biological control, 1675
balsam fir, 1760
behavior, 401
beneficial, 1083
biological control, 21
boric acid, 1362
carbaryl, 584
contour mapping, 1617
cost analysis, 1675
cotton, 805
cyfluthrin, 1362
economic threshold, 1500
elm leaf beetle, 1455
food-based bait, 584
Fraser fir, 1760
German cockroach, 1362
grasshopper, 584
host plant phenology, 1760
Hymenoptera, 1083
imidacloprid, 1455
Kaolin, 805
life table, 21
Lygus lineolaris, 401
modified hive entrances, 1647
natural enemies, 1675
oil, 584
ornamentals, 1675
particle films, 805
Pectinophora gossypiella, 805
pesticide, 1083
physical control method, 401
pink bollworm, 805
screened bottom boards, 1647
semivariogram, 1617
spatial analysis, 1617
strawberry, 401, 1500
swine production, 1362
tarnished plant bug, 1500
Trichogramma, 1083
Trogoderma variabile, 1617
vacuuming, 401
Xanthogaleruca luteola, 1455
interplanting,
associational resistance, 81
pest management, 81
Popillia japonica, 81
Rosa × *hybrida*, 81
intraspecific interspecific interactions,
Lipolexis scutellaris, 564
Lysiphlebus testaceipes, 564
Toxoptera citrida, 564
intraspecific variation,
cowpea, 489
pest management, 489
resistance, 489
seed beetle, 489
introduced basswood thrips,
host location, 777
olfactory cues, 777
pear thrips, 777
pest monitoring, 777
visual cues, 777
irradiation,
Maconellicoccus hirsutus, 1334
pink hibiscus mealybug, 1334
quarantine security, 1334
isoborneol,
bioavailability, 1267
persistence, 1267
Reticulitermes spp., 1267
soil treatments, 1267
isolation barriers,
Anastrepha fraterculus' stocks, 1237
karyotype variability, 1237
N-banding, 1237
RAPDs/PCR, 1237
ivermectin,
Hematobia irritans, 1608
sustained release bolus, 1608
ivy geranium,
generic Taylor coefficients, 482
presence-absence sampling, 482
twospotted spider mite, 482

J

Japanese beetle,
monitoring, 453
traps, 453

K

Kaolin,
cotton, 805
integrated pest management (IPM), 805
particle films, 805
Pectinophora gossypiella, 805
pink bollworm, 805
karyotype variability,
Anastrepha fraterculus' stocks, 1237
isolation barriers, 1237
N-banding, 1237
RAPDs/PCR, 1237
kdr-type resistance,
German cockroach, 885
knockdown resistance, 885
multiplex polymerase chain reaction, 885
para, 885
kernel damage,
bioassay, 220
dry weight loss, 220
host plant resistance, 220
Rhyzopertha dominica, 220
stored-products, 220
kill strips,
Anthonomus grandis, 348
boll weevil, 348
pheromone traps, 348
knockdown resistance,
German cockroach, 885
kdr-type resistance, 885
multiplex polymerase chain reaction, 885
para, 885

L
laboratory bioassays,
field insecticide tests, 1463
Rhagoletis pomonella, 1463
Laccaria bicolor,
Choristoneura occidentalis, 783
fertilization, 783
foliar nutrients, 783
larval feeding preferences, 783
minerals, 783
lady beetles,
cotton integrated pest management, 239
seasonality, 239
sticky trap, 239
trapping, 239
lambda-cyhalothrin,
acephate, 645
chlorpyrifos, 645
methomyl, 645
Solenopsis invicta, 645
toxicology, 645
Lambdina fiscellaria,
egg sampling, 768
light trap, 768
monitoring, 768
outbreak detection, 768
oviposition trap, 768
larval feeding preferences,
Choristoneura occidentalis, 783
fertilization, 783
foliar nutrients, 783
Laccaria bicolor, 783
minerals, 783
larval instar determination,
corn rootworm, 1153
Diabrotica barberi, *1153*
Diabrotica virgifera virgifera, 1153
larval sampling, 1153
larval monitoring,
funnel trap, 835
root weevil, 835
larval movement,
Diabrotica virgifera virgifera, 599
resistance management, 599
western corn rootworm, 599
larval sampling,
corn rootworm, 1153
Diabrotica barberi, *1153*
Diabrotica virgifera virgifera, 1153
larval instar determination, 1153
LdVssc1,
esfenvalerate, 1021
insect genetics, 1021
insecticide resistance, 1021
Para, 1021
leafhopper, 1221, 1369
Endopiza viteana, 1221
Maladera castanea, 1221
Pierce's disease, 1369
sharpshooter, 1369
vector, 1369
vineyards, 1221
Xylella fastidiosa, 1369
LED-plastic cup trap,
Bemisia tabaci, 543
Encarsia, 543
Eretmocerus, 543
yellow sticky card trap, 543
lepidopterous pests,
Cry1Ac, 755
Cry2Ab, 755
transgenic *Bacillus thuringiensis* cotton, 755
lesser mealworm,
Alphitobius diaperinus, 1352
house fly, 1352
Musca domestica, 1352
poultry pest management, 1352
lethal sublethal effects,
neem, 35
Neodiprion abietis, 35
lettuce,
Controlled atmosphere, 1100
postharvest pest control, 1100
postharvest quality, 1100
vacuum, 1100
life table,
biological control, 21
integrated pest management (IPM), 21
life table parameters,
Eoreuma loftini, 950
Galanthus nivalis agglutinin, 950
Galanthus nivalis L., 950
host plant resistance, 950
light modulation,
Diuraphis noxia, 194
photosynthesis, 194
plant physiology, 194
Russian wheat aphid, 194
wheat, 194
light trap,
egg sampling, 768
Lambdina fiscellaria, 768
monitoring, 768
outbreak detection, 768
oviposition trap, 768
lignin,
Anagrapha falcifera nuclear polyhedrovirus, 1066
microencapsulation, 1066
shelf-life, 1066
spray dry, 1066
Linepithema humile, 844, 850, 871
Argentine ant, 844, 871
aromatic cedar mulch, 850
control, 844
cultural control, 850
field study, 850
foraging, 871
foraging range, 844
granular bait, 871
nesting sites, 850
resurgence, 844
Lipolexis scutellaris,
intraspecific interspecific interactions, 564
Lysiphlebus testaceipes, 564
Toxoptera citrida, 564
Lolium perenne,
Festuca arundinacea, 1898
Poa pratensis, 1898
Rhizotrogus majalis, 1898
tolerance, 1898
lowbush blueberry,
gall, 1245
Hemadas nubilipennis, 1245
Vaccinium angustifolium, 1245
Luria Bertani,
Bacillus thuringiensis subsp. *israelensis*, 1039
Bengal gram, 1039
common sugar, 1039
mosquitoes, 1039
potato, 1039
Lygus lineolaris, 401, 433, 902
behavior, 401
boll weevil eradication, 902
integrated pest management, 401
malathion, 902
physical control method, 401
resistant cultivars, 433
strawberry, 401, 433
tarnished plant bug, 433, 902
vacuuming, 401
Lymantria dispar L.,
costs, 1448
spread, 1448
Lymantria monacha,
development, 43
host plant suitability, 43
survival, 43
Lysiphlebus testaceipes, 564, 975
cereal aphids, 975
intraspecific interspecific interactions, 564
Lipolexis scutellaris, 564
natural enemy threshold, 975
sequential sampling, 975
Toxoptera citrida, 564

M
Maconellicoccus hirsutus,
irradiation, 1334
pink hibiscus mealybug, 1334
quarantine security, 1334
maize, 649, 941
baculovirus, 649
crop phenology, 649
natural enemies, 649
Ostrinia nubilalis, 941
resistance management, 941
synthetic pesticides, 649
transgenic, 941
maize weevil,
control, 510
diatomaceous earth, 510
rice weevil, 510
Sitophilus, 510
stored grain, 510
Maladera castanea, 1076, 1221
Endopiza viteana, 1221
entomopathogenic nematodes, 1076
Heterorhabditis, 1076
insecticides, 1076
leafhoppers, 1221
Steinernema, 1076
vineyards, 1221
malathion, 311, 902
bait, 311
boll weevil, 311
boll weevil eradication, 902
Lygus lineolaris, 902
tarnished plant bug, 902
toxicity, 311
Mamestra brassicae,
Brassica oleracea, 143
host plant resistance, 143
Pieris brassicae, 143
Pieris rapae, 143
mark-recapture,
Diabrotica virgifera, 1750
rubidium, 1750
western corn rootworm, 1750
Zea mays, 1750
mating competitiveness,
Ceratitis capitata, 1132
sterile insect technique, 1132
mating delay,
age, 592
Cnephasia jactatana, 592
reproductive potential, 592
Tortricidae, 592
mating disruption,
apple, 58
Grapholita molesta, 58
microencapsulation, 58
sprayable pheromone, 58
maturation,
eggs, 341
oogenesis, 341
sugar, 341
yeast hydrolysate, 341
Mayetiola destructor (Say),
hydrogen phosphide fumigation, 1340
pressure, 1340
McPhail Trap,
citrus pests, 1732
fruit fly lures, 1732
nontarget insects, 1732
Tephritidae, 1732
mechanical trap,
Brahman breeding, 892
Hematobia irritans irritans, 892
IGR-bolus, 892
innate fly resistance, 892
pasture rotation, 892
mechanisms of resistance,
Diuraphis noxia, 214
wheat, 214
Medicago sativa,
Empoasca fabae, 1184
strip harvesting, 1184
Mediterranean fruit fly,
Ceratitis capitata, 662
metamorphosis, 662
phloxine B, 662
photoinsecticides, 662
Melanoplus sanguinipes, 1045, 1193
above- below-ground growth, 1193
biological control, 1045
grasshopper density, 1193
mycoinsecticide, 1045
nontarget organism, 1045
oilseed crop, 1193
predators, 1045
small-grains, 1193
stored product insects, 1045
melon fly,
cue-lure, 1780
methyl eugenol, 1780
oriental fruit fly, 1780
spinosad, 1780
metamorphosis,
Ceratitis capitata, 662
Mediterranean fruit fly, 662
phloxine B, 662
photoinsecticides, 662
Metarhizium anisopliae,
Apis mellifera, 1091
application methods, 1091
biological control, 1091
Varroa destructor, 1091
methomyl,
acephate, 645
chlorpyrifos, 645
lambda-cyhalothrin, 645
Solenopsis invicta, 645
toxicology, 645
methoprene,
cyromazine, 1300
diflubenzuron, 1300
insect growth regulator (IGR), 1300
selection, 1300
methoxychlor,
bioassay, 1016
insecticide resistance, 1016
Musca domestica, 1016
permethrin, 1016
stirofos, 1016
methoxyfenozide,
Cydia pomonella, 623
orientation tube, 623
sex pheromone, 623
wind tunnel, 623
methyl bromide alternative,
heat tolerance, 1345
heat treatment, 1345
Tribolium castaneum, 1345
methyl eugenol,
cue-lure, 1780
melon fly, 1780
oriental fruit fly, 1780
spinosad, 1780
methyl parathion,
Diabrotica virgifera virgifera, 1855
esterase, 1855
insecticide resistance, 1855
organophosphate, 1855
microbial control,
azadirachtin, 25
Bemisia, 25
Paecilomyces, 25
microencapsulation, 58, 1066
Anagrapha falcifera nuclear polyhedrovirus, 1066
apple, 58
Grapholita molesta, 58
lignin, 1066
mating disruption, 58
shelf-life, 1066
spray dry, 1066
sprayable pheromone, 58
microsatellites,
bioinvasion, 1711
Ceratitis capitata, 1711
mtDNA, 1711
polymerase chain reaction-restriction fragment length polymorphism, 1711
microsporidia,
chronic, 12
Diatraea grandiosella, 12
sublethal, 12
vertical transmission, 12
migration,
diapause, 1433
pest management, 1433

phenology, 1433
resistance management, 1433
milky disease,
Ataenius spretulus, 792
carabids, 792
golf course, 792
Paenibacillus sp., 792
staphylinids, 792
millet,
European corn borer, 361
Ostrinia nubilalis, 361
refuge, 361
trap crop, 361
minerals,
Choristoneura occidentalis, 783
fertilization, 783
foliar nutrients, 783
Laccaria bicolor, 783
larval feeding preferences, 783
mitochondrial DNA, 669, 1514
corn rootworm, 669
Diabrotica, 669
molecular diagnostics, 1514
PCR-restriction fragment-length polymorphism, 1514
Reticulitermes, 1514
species diagnostics, 669
termites, 1514
mode of action,
essential oil, 1479
natural fumigant, 1479
natural insecticide, 1479
Trialeurodes vaporariorum, 1479
model,
Ceratitis capitata, 300
cold treatment, 300
modeling,
crop rotation, 1872
Diabrotica virgifera virgifera, 1872
insect resistance management, 1872
moderate dose,
Helicoverpa zea, 914
plant resistance, 914
transgenic corn, 914
modified hive entrances,
Aethina tumida, 1647
Apis mellifera, 1647
integrated pest management, 1647
screened bottom boards, 1647
molasses bait,
Citrus thrips, 629
phloxine B, 629
photoactive dye, 629
Scirtothrips citri, 629
molecular diagnostics,
mitochondrial DNA, 1514
PCR-restriction fragment-length polymorphism, 1514
Reticulitermes, 1514
termites, 1514
monitoring, 453, 768, 826, 1821
attractants, 826
Conotrachelus nenuphar, 826
Dasineura oxycoccana, 1821
egg sampling, 768
Japanese beetle, 453
Lambdina fiscellaria, 768
light trap, 768
odor, 826
outbreak detection, 768
oviposition trap, 768
trap trees, 826
traps, 453, 1821
Vaccinium spp., 1821
Monte Carlo simulation,
alfalfa, 536
alfalfa weevil, 536
bean leaf beetle, 536
soybean, 536
mortality,
Aphis pomi, 1864
green apple aphid, 1864
imidacloprid, 1864
toxicity, 1864
mosquitoes,
Bacillus thuringiensis subsp. *israelensis*, 1039
Bengal gram, 1039
common sugar, 1039
Luria Bertani, 1039
potato, 1039
mtDNA,
bioinvasion, 1711
Ceratitis capitata, 1711
microsatellites, 1711
polymerase chain reaction-restriction fragment length polymorphism, 1711
multiplex polymerase chain reaction,
German cockroach, 885
kdr-type resistance, 885
knockdown resistance, 885
para, 885
Musca domestica, 1016, 1352
Alphitobius diaperinus, 1352
bioassay, 1016
house fly, 1352
insecticide resistance, 1016
lesser mealworm, 1352
methoxychlor, 1016
permethrin, 1016
poultry pest management, 1352
stirofos, 1016
mycoinsecticide,
biological control, 1045
Melanoplus sanguinipes, 1045
nontarget organism, 1045
predators, 1045
stored product insects, 1045
Myiopharus macellus,
biological control, 706
cultural control, 706
sunflower beetle, 706
Zygogramma exclamationis, 706

N

natural enemies, 649, 996, 1675
augmentative biological control, 1675
baculovirus, 649
cost analysis, 1675
crop phenology, 649
grain elevator, 996
integrated pest management, 1675
maize, 649
ornamentals, 1675
population dynamics, 996
synthetic pesticides, 649
natural enemy threshold,
cereal aphids, 975
Lysiphlebus testaceipes, 975
sequential sampling, 975
natural fumigant,
essential oil, 1479
mode of action, 1479
natural insecticide, 1479
Trialeurodes vaporariorum, 1479
natural insecticide,
essential oil, 1479
mode of action, 1479
natural fumigant, 1479
Trialeurodes vaporariorum, 1479
naturalyte,
cocoon penetration, 1054
Hyposoter didymator, 1054
IGRs, 1054
parasitoid, 1054
pupae, 1054
N-banding,
Anastrepha fraterculus' stocks, 1237
isolation barriers, 1237
karyotype variability, 1237
RAPDs/PCR, 1237
neem, 35, 817
Azadirachta indica, 817
lethal sublethal effects, 35
Neodiprion abietis, 35
soil application, 817
systemic effects, 817
western flower thrips, 817

Neodiprion abietis,
lethal sublethal effects, 35
neem, 35
neonicotinoid,
cottony cushion scale, 1388
IGR, 1388
organophosphate, 1388
pyrethroid, 1388
nepetalactone,
catnip, 1275
Reticulitermes, 1275
soil barrier, 1275
termite, 1275
nesting sites,
aromatic cedar mulch, 850
cultural control, 850
field study, 850
Linepithema humile, 850
Nezara viridula,
Anticarsia gemmatalis, 690
Empoasca fabae, 690
genetically modified soybeans, 690
Roundup Ready soybeans, 690
Spissistilus festinus, 690
Nilaparvata lugens,
infestation, 1798
nutrient uptake, 1798
rice root, 1798
no-choice tests,
choice tests, 53
Host plant resistance, 53
Odontota dorsalis, 53
Robinia pseudoacacia L., 53
nonrepellent termiticides,
Coptotermes formosanus, 456
exposure duration, 456
nontarget effects,
biological control, 570
Eoreuma loftini, 570
fitness, 570
Galanthus nivalis agglutinin, 570
nontarget insects,
citrus pests, 1732
fruit fly lures, 1732
McPhail Trap, 1732
Tephritidae, 1732
nontarget organism,
biological control, 1045
Melanoplus sanguinipes, 1045
mycoinsecticide, 1045
predators, 1045
stored product insects, 1045
northern corn rootworm,
Diabrotica barberi Smith Lawrence, 1420
Diabrotica virgifera virgifera LeConte, 1420
sampling efficiency, 1420
western corn rootworm, 1420
nucleopolyhedrovirus,
insecticidal activity, 1668
optical brightener, 1668
Spodoptera exigua, 1668
synergism, 1668
nutrient uptake,
infestation, 1798
Nilaparvata lugens, 1798
rice root, 1798
nutrition, 106, 370
biological control, 370
Coptotermes formosanus, 106
incipient colonies, 106
insecta, 370
Phoracantha recurva, 370
Phoracantha semipunctata, 370
pollen, 370
reproduction, 106
wood species, 106
Nysius huttoni,
development, 1119
reproduction, 1119
survival, 1119
temperature, 1119

O

Ochlerotatus,
Aedes, 1786
Bacillus thuringiensis variety *israelensis*, 1786
biocontrol, 1786
Culex, 1786
Odontota dorsalis,
choice tests, 53
Host plant resistance, 53
no-choice tests, 53
Robinia pseudoacacia L., 53
odor,
attractants, 826
Conotrachelus nenuphar, 826
monitoring, 826
trap trees, 826
oil,
carbaryl, 584
food-based bait, 584
grasshopper, 584
integrated pest management (IPM), 584
oilseed crop,
above- below-ground growth, 1193
grasshopper density, 1193
Melanoplus sanguinipes, 1193
small-grains, 1193
olfactory cues,
host location, 777
introduced basswood thrips, 777
pear thrips, 777
pest monitoring, 777
visual cues, 777
onion, 1510, 1843
Allium cepa, 1510
honey bee behavior, 1510
honey bees, 1510
onion thrips, 1843
pollination, 1510
pyrethroid, 1843
resistance, 1843
Thrips tabaci, 1843
onion thrips,
onion, 1843
pyrethroid, 1843
resistance, 1843
Thrips tabaci, 1843
oogenesis,
eggs, 341
maturation, 341
sugar, 341
yeast hydrolysate, 341
optical brightener,
insecticidal activity, 1668
nucleopolyhedrovirus, 1668
Spodoptera exigua, 1668
synergism, 1668
orchard pest management,
Bacillus thuringiensis, 879
Choristoneura rosaceana, 879
insecticide resistance, 879
Pandemis limitata, 879
spinosad, 879
organophosphate, 1388, 1855
cottony cushion scale, 1388
Diabrotica virgifera virgifera, 1855
esterase, 1855
IGR, 1388
insecticide resistance, 1855
methyl parathion, 1855
neonicotinoid, 1388
pyrethroid, 1388
organosilicones,
Aphis gossypii, 246
Frankliniella occidentalis, 246
Platynota stultana, 246
Pseudococcus maritimus, 246
Tetranychus pacificus, 246
Orgyia pseudotsugata,
Bacillus thurengiensis var. *kurstaki*, 396
diflubenzuron, 396
tebufenozide, 396
oriental fruit fly,
cue-lure, 1780
melon fly, 1780
methyl eugenol, 1780
spinosad, 1780
orientation tube,
Cydia pomonella, 623
methoxyfenozide, 623
sex pheromone, 623
wind tunnel, 623

Orius,
Carabidae, 1805
corn, 1805
imidacloprid, 1805
predator, 1805
Orius insidiosus,
Frankliniella, 1201
Geocoris punctipes, 1201
spinosad, 1201
UV reflective mulch, 1201
ornamentals,
augmentative biological control, 1675
cost analysis, 1675
integrated pest management, 1675
natural enemies, 1675
Ostrinia furnacalis,
Bt corn, 935
host plant resistance, 935
transgenic plants, 935
Ostrinia nubilalis, 361, 413, 941, 1653
Bacillus thuringiensis, 1653
Beauveria bassiana, 1653
European corn borer, 361
Helicoverpa zea, 1653
insecta, 413
maize, 941
millet, 361
refuge, 361
resistance management, 941
sweet corn, 413, 1653
transgenic, 941
trap crop, 361
Zea mays, 413
outbreak detection,
egg sampling, 768
Lambdina fiscellaria, 768
light trap, 768
monitoring, 768
oviposition trap, 768
ovicide,
contact toxicity, 990
development inhibition, 990
d-Limonene, 990
feeding deterrent, 990
fumigant toxicity, 990
oviposition behavior,
Anastrepha fraterculus, 1693
citrus, 1693
host status, 1693
pest status, 1693
oviposition trap,
egg sampling, 768
Lambdina fiscellaria, 768
light trap, 768
monitoring, 768
outbreak detection, 768

P

Paecilomyces,
azadirachtin, 25
Bemisia, 25
microbial control, 25
Paenibacillus sp.,
Ataenius spretulus, 792
carabids, 792
golf course, 792
milky disease, 792
staphylinids, 792
Pakistan,
Aphis gossypii, 875
insecticide resistance, 875
pyrethroids, 875
Pandemis limitata,
Bacillus thuringiensis, 879
Choristoneura rosaceana, 879
insecticide resistance, 879
orchard pest management, 879
spinosad, 879
Pangaeus bilineatus,
Arachis hypogaea, 1142
Cyrtomenus ciliatus, 1142
economic injury, 1142
Sehirus cinctus, 1142
Para, 885, 1021
esfenvalerate, 1021
German cockroach, 885
insect genetics, 1021
insecticide resistance, 1021
kdr-type resistance, 885
knockdown resistance, 885
LdVssc1, 1021
multiplex polymerase chain reaction, 885
parasitism,
Cirrospilus, 577
citrus, 577
dynamics, 577
peelminer, 577
parasitoid, 446, 1054
biological control, 446
biorational pesticide use, 446
cocoon penetration, 1054
Hyposoter didymator, 1054
IGRs, 1054
naturalyte, 1054
pupae, 1054
scale insects, 446
systemic insecticides, 446
particle films,
cotton, 805
integrated pest management (IPM), 805
Kaolin, 805
Pectinophora gossypiella, 805
pink bollworm, 805
pasture rotation,
Brahman breeding, 892
Hematobia irritans irritans, 892
IGR-bolus, 892
innate fly resistance, 892
mechanical trap, 892
PCR-restriction fragment-length polymorphism,
mitochondrial DNA, 1514
molecular diagnostics, 1514
Reticulitermes, 1514
termites, 1514
pear thrips,
host location, 777
introduced basswood thrips, 777
olfactory cues, 777
pest monitoring, 777
visual cues, 777
peas,
botanical insecticides, 1005
Cryptolestes, 1005
Sitophilus, 1005
traps, 1005
Tribolium, 1005
Pectinophora gossypiella,
cotton, 805
integrated pest management (IPM), 805
Kaolin, 805
particle films, 805
pink bollworm, 805
peelminer,
Cirrospilus, 577
citrus, 577
dynamics, 577
parasitism, 577
perimeter,
Tephritidae, 420
trap crop, 420
Zonosemata electa, 420
perimeter treatment,
ant exclusion, 98
granular contact insecticide, 98
repellency, 98
Perkinsiella saccharicida,
cultivar preference, 148
Fiji disease virus, 148
planthopper vector, 148
resistance, 148
sugar cane, 148
permethrin, 957, 1016
Bacillus thuringiensis corn, 957
bioassay, 1016
European corn borer, 957
insecticide resistance, 1016
methoxychlor, 1016
Musca domestica, 1016
South Dakota, 957
stirofos, 1016
univoltine ecotype, 957
yield, 957
persistence,
bioavailability, 1267
isoborneol, 1267

Reticulitermes spp., 1267
soil treatments, 1267
pest management, 81, 489, 1433
associational resistance, 81
cowpea, 489
diapause, 1433
interplanting, 81
intraspecific variation, 489
migration, 1433
phenology, 1433
Popillia japonica, 81
resistance, 489
resistance management, 1433
Rosa × *hybrida*, 81
seed beetle, 489
pest monitoring,
host location, 777
introduced basswood thrips, 777
olfactory cues, 777
pear thrips, 777
visual cues, 777
pest status,
Anastrepha fraterculus, 1693
citrus, 1693
host status, 1693
oviposition behavior, 1693
pesticide,
beneficial, 1083
Hymenoptera, 1083
integrated pest management, 1083
Trichogramma, 1083
pests,
egg predators, 71
insecticide, 71
Zea mays, 71
pet food,
community ecology, 1958
stored-product insects, 1958
trapping, 1958
Phaseolus vulgaris, 1160, 1577
antibiosis, 1577
antixenosis, 1577
economic threshold, 1160
EIL, 1160
Epilachna varivestis, 1160
host plant resistance, 1577
Thrips palmi, 1577
phenolic gylcosides,
Gypchek, 1662
gypsy moth, 1662
Populus, 1662
salicin, 1662
phenology, 1433, 1814
diapause, 1433
migration, 1433
pest management, 1433
resistance management, 1433
sampling, 1814
sticky trap, 1814
strawberry, 1814
tarnished plant bug, 1814
pheromone traps,
Anthonomus grandis, 348
boll weevil, 348
kill strips, 348
phloxine B, 629, 662
Ceratitis capitata, 662
Citrus thrips, 629
Mediterranean fruit fly, 662
metamorphosis, 662
molasses bait, 629
photoactive dye, 629
photoinsecticides, 662
Scirtothrips citri, 629
Phoracantha recurva,
biological control, 370
insecta, 370
nutrition, 370
Phoracantha semipunctata, 370
pollen, 370
Phoracantha semipunctata,
biological control, 370
insecta, 370
nutrition, 370
Phoracantha recurva, 370
pollen, 370
photoactive dye,
Citrus thrips, 629
molasses bait, 629
phloxine B, 629
Scirtothrips citri, 629
photoinsecticides,
Ceratitis capitata, 662
Mediterranean fruit fly, 662
metamorphosis, 662
phloxine B, 662
photosynthesis, 188, 194, 1168
Aphis glycines, 188
Calomycterus setarius, 1168
Cerotoma trifurcata, 1168
Diuraphis noxia, 194
fluorescence, 188
gas exchange, 188
insect-plant relations, 188
light modulation, 194
plant physiology, 194
Russian wheat aphid, 194
soybean, 1168
wheat, 194
physical control method,
behavior, 401
integrated pest management, 401
Lygus lineolaris, 401
strawberry, 401
vacuuming, 401
phytosanitary treatment,
Conotrachelus nenuphar, 1399
disinfestation, 1399
gamma radiation, 1399
Picea sp.,
AK (south-central), 31
antiaggregation, 31
Bark beetles, 31
Dendroctonus rufipennis, 31
semiochemicals, 31
Pierce's disease, 264, 1369
glassy-winged sharpshooter, 264
leafhopper, 1369
sharpshooter, 1369
vector, 264, 1369
Xylella fastidiosa, 1369
xylem, 264
Pieris brassicae,
Brassica oleracea, 143
host plant resistance, 143
Mamestra brassicae, 143
Pieris rapae, 143
Pieris rapae,
Brassica oleracea, 143
host plant resistance, 143
Mamestra brassicae, 143
Pieris brassicae, 143
pink bollworm,
cotton, 805
integrated pest management (IPM), 805
Kaolin, 805
particle films, 805
Pectinophora gossypiella, 805
pink hibiscus mealybug,
irradiation, 1334
Maconellicoccus hirsutus, 1334
quarantine security, 1334
pith,
rind, 1563
Sesamia nonagrioides, 1563
stalk strength, 1563
Zea mays, 1563
placement,
baits, 1520
Reticulitermes flavipes, 1520
Sentricon, 1520
subterranean termite control, 1520
plant compensation,
Anthonomus musculus, 407
cranberry, 407
cranberry weevil, 407
Vaccinium, 407
yield, 407
plant physiology,
Diuraphis noxia, 194
light modulation, 194
photosynthesis, 194
Russian wheat aphid, 194
wheat, 194
plant resistance, 475, 914, 1329, 1905, 1915, 1942
Aceria tosichella, 1329
Aegilops tauschii, 1329

Blissus occiduus, 1942
Buchloë dactyloides, 1942
carotenoid, 475
cassava, 1905, 1915
chlorophyll, 475
cyanogenic potential, 1905, 1915
Cyrtomenus bergi, 1905, 1915
Diuraphis noxia, 475
Helicoverpa zea, 914
moderate dose, 914
Russian wheat aphid, 475
secondary plant metabolite, 1905, 1915
transgenic corn, 914
Triticum aestivum, 1329
wheat, 475
planthopper vector,
cultivar preference, 148
Fiji disease virus, 148
Perkinsiella saccharicida, 148
resistance, 148
sugar cane, 148
Platynota stultana,
Aphis gossypii, 246
Frankliniella occidentalis, 246
organosilicones, 246
Pseudococcus maritimus, 246
Tetranychus pacificus, 246
Plodia interpunctella,
European corn borer, 503
imidacloprid, 503
Indianmeal moth, 503
thiamethoxam, 503
Plutella xylostella,
λ-cyhalothrin, 1230
indoxacarb, 1230
reduced-risk insecticide, 1230
Trichoplusia ni, 1230
Poa pratensis,
Festuca arundinacea, 1898
Lolium perenne, 1898
Rhizotrogus majalis, 1898
tolerance, 1898
pollen,
biological control, 370
insecta, 370
nutrition, 370
Phoracantha recurva, 370
Phoracantha semipunctata, 370
pollination, 547, 1215, 1510
Allium cepa, 1510
Apis mellifera, 547, 1215
bee density, 1215
colony management, 547
greenhouse, 547
honey bee, 547
honey bee behavior, 1510
honey bees, 1510
onion, 1510
rabbiteye blueberry, 1215
tomatoes, 547
Vaccinium ashei, 1215
polymerase chain reaction-restriction fragment length polymorphism,
bioinvasion, 1711
Ceratitis capitata, 1711
microsatellites, 1711
mtDNA, 1711
polyol esters,
acyl sugar, 636
biopesticide, 636
biorational insecticide, 636
sucrose octanoate, 636
surfactant, 636
Popillia japonica, 81, 1412
associational resistance, 81
conservation biological control, 1412
host location, 1412
imidacloprid, 1412
interplanting, 81
pest management, 81
Rosa × *hybrida*, 81
sublethal effects, 1412
population dynamics, 996, 1530
Diabrotica virgifera virgifera LeConte, 1530
grain elevator, 996
insecticidal traits, 1530
natural enemies, 996
resistance management, 1530
Populus,
Gypchek, 1662
gypsy moth, 1662
phenolic gylcosides, 1662
salicin, 1662
postharvest,
dried fruits, 519
Heat treatments, 519
Indianmeal moth, 519
radio frequency, 519
tree nuts, 519
postharvest disinfestation,
Callosobruchus maculatus, 1952
cold storage, 1952
cowpea weevil, 1952
dried beans, 1952
postharvest pest control,
Controlled atmosphere, 1100
lettuce, 1100
postharvest quality, 1100
vacuum, 1100
postharvest quality,
Controlled atmosphere, 1100
lettuce, 1100
postharvest pest control, 1100
vacuum, 1100
potato,
Bacillus thuringiensis subsp. *israelensis*, 1039
Bengal gram, 1039
common sugar, 1039
Luria Bertani, 1039
mosquitoes, 1039
poultry,
Alphitobius diaperinus, 1602
Beauveria bassiana, 1602
Dermestes maculatus, 1602
poultry house,
Alphitobius diaperinus, 126
control, 126
field trial, 126
insect growth regulator, 126
pyrethroid, 126
poultry pest management,
Alphitobius diaperinus, 1352
house fly, 1352
lesser mealworm, 1352
Musca domestica, 1352
precision placement,
aldicarb, 748
Cotton, 748
Frankliniella fusca, 748
Temik, 748
tobacco thrips, 748
predator, 1045, 1805
biological control, 1045
Carabidae, 1805
corn, 1805
imidacloprid, 1805
Melanoplus sanguinipes, 1045
mycoinsecticide, 1045
nontarget organism, 1045
Orius, 1805
stored product insects, 1045
presence-absence sampling,
generic Taylor coefficients, 482
ivy geranium, 482
twospotted spider mite, 482
pressure,
hydrogen phosphide fumigation, 1340
Mayetiola destructor (Say), 1340
probit analysis,
bioassay, 983
extra binomial variation, 983
generalized linear mixed model, 983
spray oils, 983
protein hydrolysates,
Bactrocera cucurbitae, 231
concentration., 231
food attractants, 231
Pseudococcus maritimus,
Aphis gossypii, 246
Frankliniella occidentalis, 246
organosilicones, 246
Platynota stultana, 246
Tetranychus pacificus, 246
Pseudoscymnus tsugae,
Adelges tsugae, 763
biological control, 763

hemlock, 763
hemlock woolly adelgid, 763
Scymnus ningshanensis, 763
Pseudotsuga menziesii,
bioassay, 377
budburst phenology, 377
Choristoneura occidentalis, 377
defoliation, 377
resistance mechanisms, 377
psocids,
beetles, 525
chlorpyrifos-methyl, 525
sorghum, 525
synergized bifenthrin, 525
pupae,
cocoon penetration, 1054
Hyposoter didymator, 1054
IGRs, 1054
naturalyte, 1054
parasitoid, 1054
pyrethroid, 126, 1388, 1843
Alphitobius diaperinus, 126
control, 126
cottony cushion scale, 1388
field trial, 126
IGR, 1388
insect growth regulator, 126
neonicotinoid, 1388
onion, 1843
onion thrips, 1843
organophosphate, 1388
poultry house, 126
resistance, 1843
Thrips tabaci, 1843
pyrethroids,
Aphis gossypii, 875
insecticide resistance, 875
Pakistan, 875
pyriproxyfen,
Bemisia, 1307
insect growth regulator (IGR), 1307
resistance, 1307
whitefly, 1307

Q

quality control,
Ceratitis capitata, 615
dominant lethal mutation, 615
sterile insect technique, 615
tsl, 615
quarantine security,
irradiation, 1334
Maconellicoccus hirsutus, 1334
pink hibiscus mealybug, 1334

R

R. flavipes,
C. formosanus, 88
foraging, 88
fractal analysis, 88
tunnel geometry, 88
rabbiteye blueberry,
Apis mellifera, 1215
bee density, 1215
pollination, 1215
Vaccinium ashei, 1215
radio frequency,
dried fruits, 519
Heat treatments, 519
Indianmeal moth, 519
postharvest, 519
tree nuts, 519
RAPDs/PCR,
Anastrepha fraterculus' stocks, 1237
isolation barriers, 1237
karyotype variability, 1237
N-banding, 1237
reciprocal cross,
Diabrotica, 131
esterase, 131
inheritance, 131
western corn rootworm, 131
reduced-risk insecticide,
λ-cyhalothrin, 1230
indoxacarb, 1230
Plutella xylostella, 1230
Trichoplusia ni, 1230
refuge, 361, 1031
Bacillus thuringiensis, 1031
Bt crops, 1031
European corn borer, 361
genetically modified crops, 1031
millet, 361
Ostrinia nubilalis, 361
resistance, 1031
trap crop, 361
remote sensing,
Coleoptera, 1770
Scarabaeidae, 1770
sound, 1770
turfgrass, 1770
repellency, 98, 461
ant exclusion, 98
Coptotermes formosanus Shiraki, 461
fipronil, 461
granular contact insecticide, 98
horizontal transmission, 461
perimeter treatment, 98
toxicity, 461
replication,
Bacillus thuringiensis, 388
efficacy, 388
experimental design, 388
spruce budworm, 388
reproduction, 106, 1119, 1612
beef cattle, 1612
Coptotermes formosanus, 106
development, 1119
heifers, 1612
Hematobia irritans, 1612
incipient colonies, 106
nutrition, 106
Nysius huttoni, 1119
survival, 1119
temperature, 1119
weight gain, 1612
wood species, 106
reproductive potential,
age, 592
Cnephasia jactatana, 592
mating delay, 592
Tortricidae, 592
resistance, 148, 489, 1031, 1290, 1307, 1843, 1886
Arachis spp., 1886
Bacillus thuringiensis, 1031, 1290
Bemisia, 1307
Bt crops, 1031
cowpea, 489
cultivar preference, 148
Fiji disease virus, 148
fitness, 1290
genetically modified crops, 1031
groundnut, 1886
Helicoverpa, 1886
Helicoverpa armigera, 1290
insect growth regulator (IGR), 1307
intraspecific variation, 489
onion, 1843
onion thrips, 1843
Perkinsiella saccharicida, 148
pest management, 489
planthopper vector, 148
pyrethroid, 1843
pyriproxyfen, 1307
refuge, 1031
seed beetle, 489
Spodoptera, 1886
sugar cane, 148
Thrips tabaci, 1843
transgenic cotton, 1290
whitefly, 1307
resistance in natural enemies,
arrhenotokous, 1626
dosage compensation, 1626
haplodiploid resistance, 1626
sexual dimorphism resistance, 1626
resistance management, 156, 173, 599, 925, 931, 941, 1433, 1530
adult emergence, 925
Bacillus thuringiensis, 156
Bacillus thuringiensis, 173
cannibalism, 931
computer simulation, 156, 173
Diabrotica virgifera virgifera, 599

Diabrotica virgifera virgifera LeConte, 1530
diapause, 1433
Helicoverpa zea, 925, 931
insecticidal traits, 1530
larval movement, 599
maize, 941
migration, 1433
Ostrinia nubilalis, 941
pest management, 1433
phenology, 1433
population dynamics, 1530
transgenic, 941
transgenic corn, 925, 931
transgenic crops, 156, 173
western corn rootworm, 599
resistance mechanisms,
bioassay, 377
budburst phenology, 377
Choristoneura occidentalis, 377
defoliation, 377
Pseudotsuga menziesii, 377
resistance screening,
genetic interaction, 202
Schizaphis graminum, 202
synthetic hexaploid wheat, 202
resistant cultivars,
Lygus lineolaris, 433
strawberry, 433
tarnished plant bug, 433
resurgence,
Argentine ant, 844
control, 844
foraging range, 844
Linepithema humile, 844
Reticulitermes, 1275, 1514
catnip, 1275
mitochondrial DNA, 1514
molecular diagnostics, 1514
nepetalactone, 1275
PCR-restriction fragment-length polymorphism, 1514
soil barrier, 1275
termite, 1275
termites, 1514
Reticulitermes flavipes, 1520, 1526
baits, 1520
Coptotermes formosanus, 1526
deltamethrin, 1526
disodium octaborate tetrahydrate, 1526
placement, 1520
Sentricon, 1520
subterranean termite control, 1520
Reticulitermes spp.,
bioavailability, 1267
isoborneol, 1267
persistence, 1267
soil treatments, 1267
Rhagoletis pomonella,
field insecticide tests, 1463
laboratory bioassays, 1463
Rhizotrogus majalis,
Festuca arundinacea, 1898
Lolium perenne, 1898
Poa pratensis, 1898
tolerance, 1898
Rhopalosiphum padi, 1585, 1792
cereal aphids, 1585, 1792
conservation tillage, 1792
conventional tillage, 1792
Schizaphis graminum, 1585
sequential sampling, 1585
winter wheat, 1585
Rhyzopertha dominica, 220, 1594
bioassay, 220
botanicals, 1594
dry weight loss, 220
feeding-deterrence, 1594
host plant resistance, 220
kernel damage, 220
Sitophilus oryzae, 1594
stored-products, 220
toxicity, 1594
Tribolium castaneum, 1594
rice bean,
bruchid resistance, 207
Callosobruchus spp., 207
Ceratotropis, 207
Vigna spp., 207
rice root,
infestation, 1798
Nilaparvata lugens, 1798
nutrient uptake, 1798
rice weevil,
control, 510
diatomaceous earth, 510
maize weevil, 510
Sitophilus, 510
stored grain, 510
rind,
pith, 1563
Sesamia nonagrioides, 1563
stalk strength, 1563
Zea mays, 1563
Robinia pseudoacacia L.,
choice tests, 53
Host plant resistance, 53
no-choice tests, 53
Odontota dorsalis, 53
root injury,
adult emergence, 714
corn rootworms, 714
soil insecticide rates, 714
root weevil,
funnel trap, 835
larval monitoring, 835
Rosa × *hybrida*,
associational resistance, 81
interplanting, 81
pest management, 81
Popillia japonica, 81
Roundup Ready soybeans,
Anticarsia gemmatalis, 690
Empoasca fabae, 690
genetically modified soybeans, 690
Nezara viridula, 690
Spissistilus festinus, 690
rubidium,
Diabrotica virgifera, 1750
mark-recapture, 1750
western corn rootworm, 1750
Zea mays, 1750
Russian wheat aphid, 194, 475, 673
carotenoid, 475
chlorophyll, 475
Diuraphis noxia, 194, 475, 673
host plant resistance, 673
light modulation, 194
photosynthesis, 194
plant physiology, 194
plant resistance, 475
wheat, 194, 475, 673

S

salicin,
Gypchek, 1662
gypsy moth, 1662
phenolic glycosides, 1662
Populus, 1662
sampling,
phenology, 1814
sticky trap, 1814
strawberry, 1814
tarnished plant bug, 1814
sampling efficiency,
Diabrotica barberi Smith Lawrence, 1420
Diabrotica virgifera virgifera LeConte, 1420
northern corn rootworm, 1420
western corn rootworm, 1420
scale insects,
biological control, 446
biorational pesticide use, 446
parasitoids, 446
systemic insecticides, 446
Scarabaeidae, 1704, 1770
Coleoptera, 1704, 1770
remote sensing, 1770
soil insects, 1704
sound, 1704, 1770
turfgrass, 1704, 1770
Schizaphis graminum, 202, 1571, 1585, 1922
Aegilops tauschii, 1571
antibiosis, 1571, 1922
antixenosis, 1922
cereal aphids, 1585
genetic interaction, 202
resistance screening, 202

Rhopalosiphum padi, 1585
sequential sampling, 1585
synthetic hexaploid wheat, 202, 1922
tolerance, 1571, 1922
wheat, 1571
winter wheat, 1585
Scirtothrips citri,
Citrus thrips, 629
molasses bait, 629
phloxine B, 629
photoactive dye, 629
screened bottom boards,
Aethina tumida, 1647
Apis mellifera, 1647
integrated pest management, 1647
modified hive entrances, 1647
Scutigerella immaculata,
bait sampling, 969
garden symphylan, 969
soil ecology, 969
soil sampling, 969
symphylid, 969
Scymnus ningshanensis,
Adelges tsugae, 763
biological control, 763
hemlock, 763
hemlock woolly adelgid, 763
Pseudoscymnus tsugae, 763
Scyphophorus acupunctatus,
aggregation pheromone, 1126
Coleoptera, 1126
Curculionidae, 1126
seasonality,
cotton integrated pest management, 239
lady beetles, 239
sticky trap, 239
trapping, 239
secondary plant metabolite, 1905, 1915
cassava, 1905, 1915
cyanogenic potential, 1905, 1915
Cyrtomenus bergi, 1905, 1915
plant resistance, 1905, 1915
seed beetle,
cowpea, 489
intraspecific variation, 489
pest management, 489
resistance, 489
Sehirus cinctus,
Arachis hypogaea, 1142
Cyrtomenus ciliatus, 1142
economic injury, 1142
Pangaeus bilineatus, 1142
selection,
cyromazine, 1300
diflubenzuron, 1300
insect growth regulator (IGR), 1300
methoprene, 1300
semiochemicals,
AK (south-central), 31
antiaggregation, 31
Bark beetles, 31
Dendroctonus rufipennis, 31
Picea sp., 31
semivariogram,
contour mapping, 1617
IPM, 1617
spatial analysis, 1617
Trogoderma variabile, 1617
Sentricon,
baits, 1520
placement, 1520
Reticulitermes flavipes, 1520
subterranean termite control, 1520
sequential sampling, 975, 1585
cereal aphids, 975, 1585
Lysiphlebus testaceipes, 975
natural enemy threshold, 975
Rhopalosiphum padi, 1585
Schizaphis graminum, 1585
winter wheat, 1585
Sesamia nonagrioides,
pith, 1563
rind, 1563
stalk strength, 1563
Zea mays, 1563
sex pheromone,
Cydia pomonella, 623
methoxyfenozide, 623
orientation tube, 623
wind tunnel, 623
sexual dimorphism resistance,
arrhenotokous, 1626
dosage compensation, 1626
haplodiploid resistance, 1626
resistance in natural enemies, 1626
sharpshooter,
leafhopper, 1369
Pierce's disease, 1369
vector, 1369
Xylella fastidiosa, 1369
shelf-life,
Anagrapha falcifera nuclear polyhedrovirus, 1066
lignin, 1066
microencapsulation, 1066
spray dry, 1066
simulation,
compensation, 1174
cotton, 1174
Helicoverpa zea, 1174
tolerance, 1174
Sitophilus,
botanical insecticides, 1005
control, 510
Cryptolestes, 1005
diatomaceous earth, 510
maize weevil, 510
peas, 1005
rice weevil, 510
stored grain, 510
traps, 1005
Tribolium, 1005
Sitophilus oryzae,
botanicals, 1594
feeding-deterrence, 1594
Rhyzopertha dominica, 1594
toxicity, 1594
Tribolium castaneum, 1594
small-grains,
above- below-ground growth, 1193
grasshopper density, 1193
Melanoplus sanguinipes, 1193
oilseed crop, 1193
smoke-tree sharpshooter,
glassy-winged sharpshooter, 1849
Homalodisca coagulata, 1849
Homalodisca lacerta, 1849
insecticide resistance, 1849
soil application,
Azadirachta indica, 817
neem, 817
systemic effects, 817
western flower thrips, 817
soil barrier,
catnip, 1275
nepetalactone, 1275
Reticulitermes, 1275
termite, 1275
soil ecology,
bait sampling, 969
garden symphylan, 969
Scutigerella immaculata, 969
soil sampling, 969
symphylid, 969
soil insecticide,
Beta vulgaris, 1426
cover crop, 1426
integrated control, 1426
Tetanops myopaeformis, 1426
soil insecticide rates,
adult emergence, 714
corn rootworms, 714
root injury, 714
soil insects,
Coleoptera, 1704
Scarabaeidae, 1704
sound, 1704
turfgrass, 1704
soil sampling,
bait sampling, 969
garden symphylan, 969
Scutigerella immaculata, 969
soil ecology, 969
symphylid, 969

soil treatments,
bioavailability, 1267
isoborneol, 1267
persistence, 1267
Reticulitermes spp., 1267
Solenopsis invicta, 94, 645
acephate, 645
bait contamination, 94
chlorpyrifos, 645
lambda-cyhalothrin, 645
methomyl, 645
toxicology, 645
sorghum,
beetles, 525
chlorpyrifos-methyl, 525
psocids, 525
synergized bifenthrin, 525
sound, 1704, 1770
Coleoptera, 1704, 1770
remote sensing, 1770
Scarabaeidae, 1704, 1770
soil insects, 1704
turfgrass, 1704, 1770
South Dakota,
Bacillus thuringiensis corn, 957
European corn borer, 957
permethrin, 957
univoltine ecotype, 957
yield, 957
soybean, 536, 1168
alfalfa, 536
alfalfa weevil, 536
bean leaf beetle, 536
Calomycterus setarius, 1168
Cerotoma trifurcata, 1168
Monte Carlo simulation, 536
photosynthesis, 1168
spatial analysis,
contour mapping, 1617
IPM, 1617
semivariogram, 1617
Trogoderma variabile, 1617
species diagnostics,
corn rootworm, 669
Diabrotica, 669
mitochondrial DNA, 669
SPF,
Chronic infection, 272
disease, 272
technique, 272
spinosad, 879, 1201, 1485, 1780, 1967
Bacillus thuringiensis, 879
Bactrocera cucurbitae, 1485
bait sprays, 1485
Choristoneura rosaceana, 879
contact toxicity, 1967
cue-lure, 1780
Frankliniella, 1201
Geocoris punctipes, 1201
GF-120 fruit fly bait, 1485
insecticide resistance, 879
melon fly, 1780
methyl eugenol, 1780
orchard pest management, 879
oriental fruit fly, 1780
Orius insidiosus, 1201
Pandemis limitata, 879
stored-product insects, 1967
surface treatments, 1967
UV reflective mulch, 1201
Spissistilus festinus,
Anticarsia gemmatalis, 690
Empoasca fabae, 690
genetically modified soybeans, 690
Nezara viridula, 690
Roundup Ready soybeans, 690
Spodoptera,
Arachis spp., 1886
groundnut, 1886
Helicoverpa, 1886
resistance, 1886
Spodoptera exigua,
insecticidal activity, 1668
nucleopolyhedrovirus, 1668
optical brightener, 1668
synergism, 1668
Spodoptera spp.,
Amaranthus spp., 798
callaloo, 798
Herpetogramma bipunctalis, 798
Spoladea recurvalis, 798
Spoladea recurvalis,
Amaranthus spp., 798
callaloo, 798
Herpetogramma bipunctalis, 798
Spodoptera spp., 798
spray dry, 292, 1066
Anagrapha falcifera nuclear polyhedrovirus, 1066
B. thuringiensis, 292
biopesticides, 292
formulation, 292
lignin, 1066
microencapsulation, 1066
shelf-life, 1066
suspensibility, 292
wettability, 292
spray oils,
bioassay, 983
extra binomial variation, 983
generalized linear mixed model, 983
probit analysis, 983
sprayable pheromone,
apple, 58
Grapholita molesta, 58
mating disruption, 58
microencapsulation, 58
spread,
costs, 1448
Lymantria dispar L., 1448
spruce budworm, 280, 388
Bacillus thuringiensis, 280, 388
Choristoneura fumiferana, 280
Double application, 280
efficacy, 388
experimental design, 388
replication, 388
sublethal effects, 280
square loss,
canopy structure, 730
Gossypium hirsutum, 730
Helicoverpa, 730
terminal removal, 730
yield, 730
stalk strength,
pith, 1563
rind, 1563
Sesamia nonagrioides, 1563
Zea mays, 1563
staphylinids,
Ataenius spretulus, 792
carabids, 792
golf course, 792
milky disease, 792
Paenibacillus sp., 792
Steinernema,
entomopathogenic nematodes, 1076
Heterorhabditis, 1076
insecticides, 1076
Maladera castanea, 1076
Steinernema spp.,
application methodology, 287
entomopathogenic nematodes, 287
Heterorhabditis bacteriophora, 287
Vitis, 287
sterile insect technique, 615, 1132, 1405, 1724
Ceratitis capitata, 615, 1132, 1405
Cryptophlebia leucotreta, 1724
dominant lethal mutation, 615
fecundity, 1724
fertility, 1724
gamma radiation, 1724
GF-120, 1405
mating competitiveness, 1132
quality control, 615
tsl, 615
sticky trap, 239, 1814
cotton integrated pest management, 239
lady beetles, 239
phenology, 1814
sampling, 1814
seasonality, 239
strawberry, 1814

tarnished plant bug, 1814
trapping, 239
stirofos,
bioassay, 1016
insecticide resistance, 1016
methoxychlor, 1016
Musca domestica, 1016
permethrin, 1016
stored grain,
control, 510
diatomaceous earth, 510
maize weevil, 510
rice weevil, 510
Sitophilus, 510
stored-product insects, 1045, 1958, 1967
biological control, 1045
community ecology, 1958
contact toxicity, 1967
Melanoplus sanguinipes, 1045
mycoinsecticide, 1045
nontarget organism, 1045
pet food, 1958
predators, 1045
spinosad, 1967
surface treatments, 1967
trapping, 1958
stored-products,
bioassay, 220
dry weight loss, 220
host plant resistance, 220
kernel damage, 220
Rhyzopertha dominica, 220
strawberry, 401, 433, 1500, 1814
behavior, 401
economic threshold, 1500
integrated pest management, 401, 1500
Lygus lineolaris, 401, 433
phenology, 1814
physical control method, 401
resistant cultivars, 433
sampling, 1814
sticky trap, 1814
tarnished plant bug, 433, 1500, 1814
vacuuming, 401
strip harvesting,
Empoasca fabae, 1184
Medicago sativa, 1184
sublethal,
chronic, 12
Diatraea grandiosella, 12
microsporidia, 12
vertical transmission, 12
sublethal effects, 280, 1412
Bacillus thuringiensis, 280
Choristoneura fumiferana, 280
conservation biological control, 1412
Double application, 280
host location, 1412
imidacloprid, 1412
Popillia japonica, 1412
spruce budworm, 280
subterranean termite control,
baits, 1520
placement, 1520
Reticulitermes flavipes, 1520
Sentricon, 1520
subterranean termites, 1259, 1828
American cockroach, 1259
Australia, 1828
bait technology, 1828
chitin synthesis inhibitor, 1828
chlorfluazuron, 1828
dog, 1259
drywood termites, 1259
termite detection, 1259
termite-damaged wood, 1259
sucrose octanoate,
acyl sugar, 636
biopesticide, 636
biorational insecticide, 636
polyol esters, 636
surfactant, 636
sugar,
eggs, 341
maturation, 341
oogenesis, 341
yeast hydrolysate, 341
sugarcane, 148, 1929
areawide pest management, 1929
cultivar preference, 148
Eoreuma loftini, 1929
Fiji disease virus, 148
host plant resistance, 1929
Perkinsiella saccharicida, 148
planthopper vector, 148
resistance, 148
sunflower beetle,
biological control, 706
cultural control, 706
Myiopharus macellus, 706
Zygogramma exclamationis, 706
surface treatments,
contact toxicity, 1967
spinosad, 1967
stored-product insects, 1967
surfactant,
acyl sugar, 636
biopesticide, 636
biorational insecticide, 636
polyol esters, 636
sucrose octanoate, 636
survival, 43, 1119
development, 43, 1119
host plant suitability, 43
Lymantria monacha, 43
Nysius huttoni, 1119
reproduction, 1119
temperature, 1119
susceptibility,
Bacillus thuringiensis, 1838
Chilo suppressalis, 1838
China, 1838
suspensibility,
B. thuringiensis, 292
biopesticides, 292
formulation, 292
spray drying, 292
wettability, 292
sustained release bolus,
Hematobia irritans, 1608
ivermectin, 1608
sweet corn, 413, 1653
Bacillus thuringiensis, 1653
Beauveria bassiana, 1653
Helicoverpa zea, 1653
insecta, 413
Ostrinia nubilalis, 413, 1653
Zea mays, 413
swine production,
boric acid, 1362
cyfluthrin, 1362
German cockroach, 1362
integrated pest management (IPM), 1362
symphylid,
bait sampling, 969
garden symphylan, 969
Scutigerella immaculata, 969
soil ecology, 969
soil sampling, 969
synergism,
insecticidal activity, 1668
nucleopolyhedrovirus, 1668
optical brightener, 1668
Spodoptera exigua, 1668
synergized bifenthrin,
beetles, 525
chlorpyrifos-methyl, 525
psocids, 525
sorghum, 525
synthetic hexaploid wheat, 202, 1922
antibiosis, 1922
antixenosis, 1922
genetic interaction, 202
resistance screening, 202
Schizaphis graminum, 202, 1922
tolerance, 1922
synthetic pesticides,
baculovirus, 649
crop phenology, 649
maize, 649
natural enemies, 649
systemic effects,
Azadirachta indica, 817
neem, 817
soil application, 817
western flower thrips, 817

systemic insecticides,
biological control, 446
biorational pesticide use, 446
parasitoids, 446
scale insects, 446

T

Tapinoma melanocephalum,
ants, 856
insecticides, 856
toxic baits, 856
tarnished plant bug, 433, 902, 1500, 1814
boll weevil eradication, 902
economic threshold, 1500
integrated pest management, 1500
Lygus lineolaris, 433, 902
malathion, 902
phenology, 1814
resistant cultivars, 433
sampling, 1814
sticky trap, 1814
strawberry, 433, 1500, 1814
tebufenozide,
Bacillus thurengiensis var. *kurstaki*, 396
diflubenzuron, 396
Orgyia pseudotsugata, 396
tebufenpyrad,
Apis mellifera, 259
Apistan, 259
CheckMite+, 259
honey bee, 259
varroa, 259
technique,
Chronic infection, 272
disease, 272
SPF, 272
Temik,
aldicarb, 748
Cotton, 748
Frankliniella fusca, 748
precision placement, 748
tobacco thrips, 748
temperature, 609, 1119
Chrysomelidae, 609
Coleoptera, 609
degree-days, 609
development, 1119
Nysius huttoni, 1119
reproduction, 1119
survival, 1119
Tephritidae, 334, 420, 1108, 1732
Anastrepha, 334
Anastrepha fraterculus, 1108
Argentina, 1108
attractants, 334
Ceratitis capitata, 1108
citrus pests, 1732
fruit fly lures, 1732
host plants, 1108
human urine, 334
McPhail Trap, 1732
nontarget insects, 1732
perimeter, 420
trap crop, 420
Zonosemata electa, 420
terminal removal,
canopy structure, 730
Gossypium hirsutum, 730
Helicoverpa, 730
square loss, 730
yield, 730
termite, 117, 1275, 1514
catnip, 1275
Coptotermes formosanus, 117
foam, 117
imidacloprid, 117
mitochondrial DNA, 1514
molecular diagnostics, 1514
nepetalactone, 1275
PCR-restriction fragment-length polymorphism, 1514
Reticulitermes, 1275, 1514
soil barrier, 1275
trees, 117
termite detection,
American cockroach, 1259
dog, 1259
drywood termites, 1259
subterranean termites, 1259
termite-damaged wood, 1259
termite-damaged wood,
American cockroach, 1259
dog, 1259
drywood termites, 1259
subterranean termites, 1259
termite detection, 1259
territoriality,
ants, 251
biological control, 251
hemiptera, 251
insecticides, 251
Tetanops myopaeformis,
Beta vulgaris, 1426
cover crop, 1426
integrated control, 1426
soil insecticide, 1426
Tetranychus pacificus,
Aphis gossypii, 246
Frankliniella occidentalis, 246
organosilicones, 246
Platynota stultana, 246
Pseudococcus maritimus, 246
thiamethoxam,
European corn borer, 503
imidacloprid, 503
Indianmeal moth, 503
Plodia interpunctella, 503
thrips
Frankliniella occidentalis, 1
tomato spotted wilt virus, 1
Thrips palmi,
antibiosis, 1577
antixenosis, 1577
host plant resistance, 1577
Phaseolus vulgaris, 1577
Thrips tabaci,
onion, 1843
onion thrips, 1843
pyrethroid, 1843
resistance, 1843
tissue difference,
cotton bollworm, 1283
cytochrome P450 monooxygenase, 1283
Helicoverpa armigera, 1283
induction, 1283
tobacco thrips,
aldicarb, 748
Cotton, 748
Frankliniella fusca, 748
precision placement, 748
Temik, 748
Togo,
Coffee tree, 1473
economic damage threshold, 1473
estimation of losses, 1473
Hypothenemus hampei, 1473
tolerance, 1174, 1571, 1898, 1922
Aegilops tauschii, 1571
antibiosis, 1571, 1922
antixenosis, 1922
compensation, 1174
cotton, 1174
Festuca arundinacea, 1898
Helicoverpa zea, 1174
Lolium perenne, 1898
Poa pratensis, 1898
Rhizotrogus majalis, 1898
Schizaphis graminum, 1571, 1922
simulation, 1174
synthetic hexaploid wheat, 1922
wheat, 1571
tomato spotted wilt virus
Frankliniella occidentalis, 1
thrips, 1
tomatoes,
Apis mellifera, 547
colony management, 547
greenhouse, 547
honey bee, 547
pollination, 547
Tortricidae,
age, 592
Cnephasia jactatana, 592
mating delay, 592
reproductive potential, 592
toxic baits,
ants, 856
insecticides, 856

Tapinoma melanocephalum, 856
toxicity, 311, 461, 863, 1559, 1594, 1864
Aedes albopictus, 1559
Anopheles quadrimaculatus, 1559
Aphis pomi, 1864
bait, 311
Blattella germanica, 863
boll weevil, 311
botanicals, 1594
Coptotermes formosanus Shiraki, 461
Culex nigripalpus, 1559
Ebeling choice box, 863
feeding-deterrence, 1594
field performance, 863
fipronil, 461
green apple aphid, 1864
horizontal transmission, 461
imidacloprid, 1864
indoxacarb, 863
insecticidal bait, 863
malathion, 311
mortality, 1864
repellency, 461
Rhyzopertha dominica, 1594
Sitophilus oryzae, 1594
Tribolium castaneum, 1594
toxicology,
acephate, 645
chlorpyrifos, 645
lambda-cyhalothrin, 645
methomyl, 645
Solenopsis invicta, 645
Toxoptera citrida,
intraspecific interspecific interactions, 564
Lipolexis scutellaris, 564
Lysiphlebus testaceipes, 564
transgenic,
maize, 941
Ostrinia nubilalis, 941
resistance management, 941
transgenic *Bacillus thuringiensis* cotton,
Cry1Ac, 755
Cry2Ab, 755
lepidopterous pests, 755
transgenic corn, 914, 925, 931
adult emergence, 925
cannibalism, 931
Helicoverpa zea, 914, 925, 931
moderate dose, 914
plant resistance, 914
resistance management, 925, 931
transgenic cotton, 1290, 1322
Bacillus thuringiensis, 1290, 1322
China, 1322
fitness, 1290
Helicoverpa armigera, 1290, 1322
resistance, 1290
transgenic crops, 156, 173
Bacillus thuringiensis, 156
Bacillus thuringiensis, 173
computer simulation, 156, 173
resistance management, 156, 173
transgenic plants,
Bt corn, 935
host plant resistance, 935
Ostrinia furnacalis, 935
transgenics,
Bollgard, 699
Bollgard II, 699
economic injury, 699
trap crop, 361, 420
European corn borer, 361
millet, 361
Ostrinia nubilalis, 361
perimeter, 420
refuge, 361
Tephritidae, 420
Zonosemata electa, 420
trap trees,
attractants, 826
Conotrachelus nenuphar, 826
monitoring, 826
odor, 826
trapping, 239, 680, 1958
Anthonomus grandis, 680
boll weevil, 680
community ecology, 1958
cotton, 680
cotton integrated pest management, 239
lady beetles, 239
pet food, 1958
seasonality, 239
sticky trap, 239
stored-product insects, 1958
traps, 453, 1005, 1821
botanical insecticides, 1005
Cryptolestes, 1005
Dasineura oxycoccana, 1821
Japanese beetle, 453
monitoring, 453, 1821
peas, 1005
Sitophilus, *1005*
Tribolium, 1005
Vaccinium spp., 1821
tree nuts,
dried fruits, 519
Heat treatments, 519
Indianmeal moth, 519
postharvest, 519
radio frequency, 519
trees,
Coptotermes formosanus, 117
foam, 117
imidacloprid, 117
termite, 117
Trialeurodes vaporariorum,
essential oil, 1479
mode of action, 1479
natural fumigant, 1479
natural insecticide, 1479
Tribolium,
botanical insecticides, 1005
Cryptolestes, 1005
peas, 1005
Sitophilus, *1005*
traps, 1005
Tribolium castaneum, 1345, 1594
botanicals, 1594
feeding-deterrence, 1594
heat tolerance, 1345
heat treatment, 1345
methyl bromide alternative, 1345
Rhyzopertha dominica, 1594
Sitophilus oryzae, 1594
toxicity, 1594
Trichogramma,
beneficial, 1083
Hymenoptera, 1083
integrated pest management, 1083
pesticide, 1083
Trichoplusia ni,
λ-cyhalothrin, 1230
indoxacarb, 1230
Plutella xylostella, 1230
reduced-risk insecticide, 1230
Triticum aestivum,
Aceria tosichella, 1329
Aegilops tauschii, 1329
plant resistance, 1329
Trogoderma variabile,
contour mapping, 1617
IPM, 1617
semivariogram, 1617
spatial analysis, 1617
tsl,
Ceratitis capitata, 615
dominant lethal mutation, 615
quality control, 615
sterile insect technique, 615
tunnel geometry,
C. formosanus, 88
foraging, 88
fractal analysis, 88
R. flavipes, 88
turfgrass, 1704, 1770
Coleoptera, 1704, 1770
remote sensing, 1770
Scarabaeidae, 1704, 1770
soil insects, 1704
sound, 1704, 1770
twospotted spider mite,
generic Taylor coefficients, 482

ivy geranium, 482
presence-absence sampling, 482

U

univoltine ecotype,
Bacillus thuringiensis corn, 957
European corn borer, 957
permethrin, 957
South Dakota, 957
yield, 957
UV reflective mulch,
Frankliniella, 1201
Geocoris punctipes, 1201
Orius insidiosus, 1201
spinosad, 1201

V

Vaccinium,
Anthonomus musculus, 407
cranberry, 407
cranberry weevil, 407
plant compensation, 407
yield, 407
Vaccinium angustifolium,
gall, 1245
Hemadas nubilipennis, 1245
lowbush blueberry, 1245
Vaccinium ashei,
Apis mellifera, 1215
bee density, 1215
pollination, 1215
rabbiteye blueberry, 1215
Vaccinium spp.,
Dasineura oxycoccana, 1821
monitoring, 1821
traps, 1821
vacuum,
Controlled atmosphere, 1100
lettuce, 1100
postharvest pest control, 1100
postharvest quality, 1100
vacuuming,
behavior, 401
integrated pest management, 401
Lygus lineolaris, 401
physical control method, 401
strawberry, 401
varroa,
Apis mellifera, 259
Apistan, 259
CheckMite+, 259
honey bee, 259
tebufenpyrad, 259
Varroa destructor,
Apis mellifera, 1091
application methods, 1091
biological control, 1091
Metarhizium anisopliae, 1091
vector, 264, 1369
glassy-winged sharpshooter, 264
leafhopper, 1369
Pierce's disease, 264, 1369
sharpshooter, 1369
Xylella fastidiosa, 1369
xylem, 264
vertical transmission,
chronic, 12
Diatraea grandiosella, 12
microsporidia, 12
sublethal, 12
Vigna spp.,
bruchid resistance, 207
Callosobruchus spp., 207
Ceratotropis, 207
rice bean, 207
vineyards,
Endopiza viteana, 1221
leafhoppers, 1221
Maladera castanea, 1221
virus,
Bemisia tabaci, 322
biotype, 322
esterase, 322
visual cues,
host location, 777
introduced basswood thrips, 777
olfactory cues, 777
pear thrips, 777
pest monitoring, 777
Vitis,
application methodology, 287
entomopathogenic nematodes, 287
Heterorhabditis bacteriophora, 287
Steinernema spp., 287

W

weight gain,
beef cattle, 1612
heifers, 1612
Hematobia irritans, 1612
reproduction, 1612
West Africa,
cotton, 468
Helicoverpa armigera, 468
insecticide mixtures, 468
insecticide resistance, 468
western corn rootworm, 131, 599, 737, 1420, 1738, 1750
Chrysomelidae, 1738
crop rotation, 737
Cry3Bb1, 1738
Diabrotica, 131
Diabrotica barberi Smith Lawrence, 1420
Diabrotica virgifera, 1750
Diabrotica virgifera virgifera, 599, 737, 1738
Diabrotica virgifera virgifera LeConte, 1420
esterase, 131
inheritance, 131
insect resistance management, 1738
larval movement, 599
mark-recapture, 1750
northern corn rootworm, 1420
reciprocal cross, 131
resistance management, 599
rubidium, 1750
sampling efficiency, 1420
Zea mays, 1750
western flower thrips,
Azadirachta indica, 817
neem, 817
soil application, 817
systemic effects, 817
wettability,
B. thuringiensis, 292
biopesticides, 292
formulation, 292
spray drying, 292
suspensibility, 292
wheat, 194, 214, 352, 475, 673, 1571
Aegilops tauschii, 1571
antibiosis, 1571
carotenoid, 475
chlorophyll, 475
Diuraphis noxia, 194, 214, 352, 475, 673
host plant resistance, 352, 673
light modulation, 194
mechanisms of resistance, 214
photosynthesis, 194
plant physiology, 194
plant resistance, 475
Russian wheat aphid, 194, 475, 673
Schizaphis graminum, 1571
tolerance, 1571
whitefly,
Bemisia, 1307
insect growth regulator (IGR), 1307
pyriproxyfen, 1307
resistance, 1307
wind tunnel,
Cydia pomonella, 623
methoxyfenozide, 623
orientation tube, 623
sex pheromone, 623
winter wheat,
cereal aphids, 1585
Rhopalosiphum padi, 1585
Schizaphis graminum, 1585
sequential sampling, 1585

wood species,
Coptotermes formosanus, 106
incipient colonies, 106
nutrition, 106
reproduction, 106

X

Xanthogaleruca luteola,
abamectin, 1455
elm leaf beetle, 1455
imidacloprid, 1455
integrated pest management, 1455
Xylella fastidiosa,
leafhopper, 1369
Pierce's disease, 1369
sharpshooter, 1369
vector, 1369
xylem,
glassy-winged sharpshooter, 264
Pierce's disease, 264
vector, 264

Y

yeast hydrolysate,
eggs, 341
maturation, 341
oogenesis, 341
sugar, 341
yellow sticky card trap,
Bemisia tabaci, 543
Encarsia, 543
Eretmocerus, 543
LED-plastic cup trap, 543
yield, 407, 730, 957
Anthonomus musculus, 407
Bacillus thuringiensis corn, 957
canopy structure, 730
cranberry, 407
cranberry weevil, 407
European corn borer, 957
Gossypium hirsutum, 730
Helicoverpa, 730
permethrin, 957
plant compensation, 407
South Dakota, 957
square loss, 730
terminal removal, 730
univoltine ecotype, 957
Vaccinium, 407

Z

Zea mays, 71, 413, 909, 1563, 1750
corn earworm, 909
cotton bollworm, 909
cross-resistance, 909
Diabrotica virgifera, 1750
egg predators, 71
Helicoverpa, 909
host plant resistance, 909
insecta, 413
insecticide, 71
mark-recapture, 1750
Ostrinia nubilalis, 413
pests, 71
pith, 1563
rind, 1563
rubidium, 1750
Sesamia nonagrioides, 1563
stalk strength, 1563
sweet corn, 413
western corn rootworm, 1750
Zonosemata electa,
perimeter, 420
Tephritidae, 420
trap crop, 420
Zygogramma exclamationis,
biological control, 706
cultural control, 706
Myiopharus macellus, 706
sunflower beetle, 706